SANTA CRUZ BIOTECHNOLOGY, INC.

NP2 (22): sc-136183



BACKGROUND

Long pentraxins are a family of highly conserved proteins that are expressed in the brain and central nervous system, and form multimeric complexes. Neuronal pentraxin 1 (NP1), NP2 and neuronal pentraxin receptor (NPR) are members of the long pentraxins that represent a neuronal uptake pathway that may function during synapse formation and remodeling. The NP1 gene is located on chromosome 17q25.3 and the protein product mediates the uptake of synaptic material, including the presynaptic snake venom toxin, taipoxin. NP2, whose function is unknown, is located on chromosome 7q22.1 and like NP1 contains several potential N-linked glycosylation sites. NPR is expressed on the cell membrane and can form heteropentamers with NP1 and NP2 that can be released from the cell membrane by proteolysis.

REFERENCES

- Hsu, Y.C. and Perin, M.S. 1995. Human neuronal pentraxin II (NPTX2): conservation, genomic structure, and chromosomal localization. Genomics 28: 220-227.
- Dodds, D.C., Omeis, I.A., Cushman, S.J., Helms, J.A. and Perin, M.S. 1997. Neuronal pentraxin receptor, a novel putative integral membrane pentraxin that interacts with neuronal pentraxin 1 and 2 and taipoxin-associated calcium-binding protein 49. J. Biol. Chem. 272: 21488-21494.
- Kirkpatrick, L.L., Matzuk, M.M., Dodds, D.C. and Perin, M.S. 2000. Biochemical interactions of the neuronal pentraxins. Neuronal pentraxin (NP) receptor binds to taipoxin and taipoxin-associated calcium-binding protein 49 via NP1 and NP2. J. Biol. Chem. 275: 17786-17792.
- 4. Online Mendelian Inheritance in Man, OMIM™. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 600750. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- Park, J.K., Ryu, J.K., Lee, K.H., Lee, J.K., Yoon, W.J., Lee, S.H., Yoo, J.W., Woo, S.M., Lee, G.Y., Lee, C.H., Kim, Y.T. and Yoon, Y.B. 2007. Quantitative analysis of NPTX2 hypermethylation is a promising molecular diagnostic marker for pancreatic cancer. Pancreas 35: 9-15.
- Marui, T., Koishi, S., Funatogawa, I., Yamamoto, K., Matsumoto, H., Hashimoto, O., Ishijima, M., Nanba, E., Nishida, H., Sugiyama, T., Kasai, K., Watanabe, K., Kano, Y., Kato, N. and Sasaki, T. 2007. No association between the neuronal pentraxin II gene polymorphism and autism. Prog. Neuropsychopharmacol. Biol. Psychiatry 31: 940-943.
- Moran, L.B., Hickey, L., Michael, G.J., Derkacs, M., Christian, L.M., Kalaitzakis, M.E., Pearce, R.K. and Graeber, M.B. 2008. Neuronal pentraxin II is highly upregulated in Parkinson's disease and a novel component of Lewy bodies. Acta Neuropathol. 115: 471-478.

CHROMOSOMAL LOCATION

Genetic locus: NPTX2 (human) mapping to 7q22.1; Nptx2 (mouse) mapping to 5 G2.

SOURCE

NP2 (22) is a mouse monoclonal antibody raised against amino acids 137-312 of NP2 of rat origin.

PRODUCT

Each vial contains 50 $\mu g~lgG_1$ in 0.5 ml of PBS with < 0.1% sodium azide, 0.1% gelatin, 20% glycerol, and 0.04% stabilizer protein.

APPLICATIONS


NP2 (22) is recommended for detection of NP2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for NP2 siRNA (h): sc-42095, NP2 siRNA (m): sc-42096, NP2 shRNA Plasmid (h): sc-42095-SH, NP2 shRNA Plasmid (m): sc-42096-SH, NP2 shRNA (h) Lentiviral Particles: sc-42095-V and NP2 shRNA (m) Lentiviral Particles: sc-42096-V.

Molecular Weight of NP2: 47 kDa.

Positive Controls: H4 cell lysate: sc-2408 or rat brain extract: sc-2392.

DATA



NP2 (22): sc-136183. Western blot analysis of NP2 expression in rat brain tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures. Not for resale.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.