

# NHERF-1 (6): sc-136228

## BACKGROUND

The Na<sup>+</sup>/H<sup>+</sup> exchange protein (NHE3) functions in transepithelial Na<sup>+</sup> absorption and is primarily expressed in the intestinal and renal brush border membrane. NHE3 regulatory factor 1 (NHERF-1) interacts with NHE3 through two PDZ (for PSD-95, discs-large and ZO-1 homology) domains, which are protein-protein interaction modules that associate with specific carboxy-terminal motifs on target proteins. Also known as EBP50, NHERF-1 facilitates cAMP inhibition of NHE3 to decrease Na<sup>+</sup> adsorption. NHERF-1 functions as a scaffold for an essential multiprotein complex of Ezrin and NHE3 for cAMP-mediated phosphorylation and consequent inhibition of NHE3. The amino-terminal PDZ domain regulates the dimerization of NHERF-1 *in vivo*. G protein-coupled receptor kinase 6A phosphorylates NHERF-1 at Ser 289 via a PDZ domain-mediated interaction. NHERF-2, also known as E3KARP, is an ubiquitously expressed protein which also functions in NHE2 regulation.

## REFERENCES

1. Aronson, P.S., et al. 1986. Molecular-properties and physiological roles of the renal Na<sup>+</sup>/H<sup>+</sup> exchanger. *Curr. Top. Mem. Trans.* 26: 57-75.
2. Sheng, M. 1996. PDZs and receptor/channel clustering: rounding up the latest suspects. *Neuron* 17: 575-578.
3. Yun, C.H., et al. 1997. cAMP-mediated inhibition of the epithelial brush border Na<sup>+</sup>/H<sup>+</sup> exchanger, NHE3, requires an associated regulatory protein. *Proc. Natl. Acad. Sci. USA* 94: 3010-3015.
4. Poulat, F., et al. 1997. The human testis determining factor SRY binds a nuclear factor containing PDZ protein interaction domains. *J. Biol. Chem.* 272: 7167-7172.
5. Imai, K., et al. 1998. Genomic structure and sequence of a human homologue (NTHL1/NTH1) of *Escherichia coli* endonuclease III with those of the adjacent parts of TSC2 and SLC9A3R2 genes. *Gene* 222: 287-295.
6. Hall, R.A., et al. 1999. G protein-coupled receptor kinase 6A phosphorylates the Na<sup>+</sup>/H<sup>+</sup> exchanger regulatory factor via a PDZ domain-mediated interaction. *J. Biol. Chem.* 274: 24328-24334.
7. Weinman, E.J., et al. 2000. NHERF associations with sodium-hydrogen exchanger isoform 3 (NHE3) and Ezrin are essential for cAMP-mediated phosphorylation and inhibition of NHE3. *Biochemistry* 39: 6123-6129.
8. Shenolikar, S., et al. 2001. N-terminal PDZ domain is required for NHERF dimerization. *FEBS Lett.* 489: 233-236.
9. Brdicková, N., et al. 2001. Interaction between two adapter proteins, PAG and EBP50: a possible link between membrane rafts and Actin cytoskeleton. *FEBS Lett.* 507: 133-136.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: SLC9A3R1 (human) mapping to 17q25.1.

## SOURCE

NHERF-1 (6) is a mouse monoclonal antibody raised against amino acids 128-249 of NHERF-1 of human origin.

## PRODUCT

Each vial contains 50 µg IgG<sub>1</sub> in 500 µl PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-136228 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## APPLICATIONS


NHERF-1 (6) is recommended for detection of NHERF-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for NHERF-1 siRNA (h): sc-63330, NHERF-1 shRNA Plasmid (h): sc-63330-SH and NHERF-1 shRNA (h) Lentiviral Particles: sc-63330-V.

Molecular Weight of NHERF-1: 50 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409, Caki-1 cell lysate: sc-2224 or SK-N-MC cell lysate: sc-2237.

## DATA



NHERF-1 (6): sc-136228. Western blot analysis of NHERF-1 expression in human endothelial whole cell lysate.

## SELECT PRODUCT CITATIONS

1. Luo, M., et al. 2020. LncRNA LINC00483 promotes gastric cancer development through regulating MAPK1 expression by sponging miR-490-3p. *Biol. Res.* 53: 14.

## RESEARCH USE

For research use only, not for use in diagnostic procedures. Not for resale.