

# MEK-7 (40): sc-136337

## BACKGROUND

A family of protein kinases located upstream of the MAP kinases and responsible for their activation has been identified. The prototype member of this family, designated MAP kinase kinase, or MEK-1, specifically phosphorylates the MAP kinase regulatory threonine and tyrosine residues present in the Thr-Glu-Tyr motif of ERK. A second MEK family member, MEK-2, resembles MEK-1 in its substrate specificity. MEK-3 (or Mkk3) functions to activate p38 MAP kinase, and MEK-4 (also called SEK1 or Mkk4) activates both p38 and JNK MAP kinases. MEK-5 appears to specifically phosphorylate ERK 5, whereas MEK-6 phosphorylates p38 and p38b. MEK-7 (or Mkk7) phosphorylates and activates the JNK signal transduction pathway.

## REFERENCES

1. Crews, C.M., Alessandrini, A. and Erikson, R.L. 1992. The primary structure of MEK, a protein kinase that phosphorylates the ERK gene product. *Science* 258: 478-480.
2. Wu, J., Harrison, J.K., Dent, P., Lynch, K.R., Weber, M.J. and Sturgill, T.W. 1993. Identification and characterization of a new mammalian mitogen-activated protein kinase kinase, MKK-2. *Mol. Cell. Biol.* 13: 4539-4548.
3. Dérijard, B., Raingeaud, J., Barrett, T., Wu, I.H., Han, J., Ulevitch, R.J. and Davis, R.J. 1995. Independent human MAP-kinase signal transduction pathways defined by MEK and MKK isoforms. *Science* 267: 682-685.
4. Zhou, G., Bao, Z.Q. and Dixon, J.E. 1995. Components of a new human protein kinase signal transduction pathway. *J. Biol. Chem.* 270: 12665-12669.
5. Han, J., Lee, J.D., Jiang, Y., Li, Z., Feng, L. and Ulevitch, R.J. 1996. Characterization of the structure and function of a novel MAP kinase kinase (MKK-6). *J. Biol. Chem.* 271: 2886-2891.
6. Jiang, Y., Chen, C., Li, Z., Guo, W., Gegner, J.A., Lin, S. and Han, J. 1996. Characterization of the structure and function of a new mitogen-activated protein kinase (p38 $\beta$ ). *J. Biol. Chem.* 271: 17920-17926.
7. Tournier, C., Whitmarsh, A.J., Cavanagh, J., Barrett, T. and Davis, R.J. 1997. Mitogen-activated protein kinase kinase-7 is an activator of the c-Jun NH<sub>2</sub>-terminal kinase. *Proc. Natl. Acad. Sci. USA* 94: 7337-7442.
8. Holland, P.M., Suzanne, M., Campbell, J.S., Noselli, S. and Cooper, J.A. 1997. Mkk7 is a stress-activated mitogen-activated protein kinase functionally related to hemipterous. *J. Biol. Chem.* 272: 24994-24998.

## CHROMOSOMAL LOCATION

Genetic locus: Map2k7 (mouse) mapping to 8 A1.1.

## SOURCE

MEK-7 (40) is a mouse monoclonal antibody raised against amino acids 42-157 of MEK-7 of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

MEK-7 (40) is recommended for detection of MEK-7 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for MEK-7 siRNA (m): sc-35916, MEK-7 shRNA Plasmid (m): sc-35916-SH and MEK-7 shRNA (m) Lentiviral Particles: sc-35916-V.

Molecular Weight of MEK-7 isoforms: 39-59 kDa.


Positive Controls: RAW 264.7 whole cell lysate: sc-2211, rat brain extract: sc-2392 or KNRK whole cell lysate: sc-2214.

## RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



MEK-7 (40): sc-136337. Western blot analysis of MEK-7 expression in RAW 264.7 whole cell lysate.



MEK-7 (40): sc-136337. Immunofluorescence staining of methanol-fixed NIH/3T3 cells showing cytoplasmic localization.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures. Not for resale.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.