

# TFIIH p80 (19): sc-136376

## BACKGROUND

Initiation of transcription from protein-coding genes in eukaryotes is a complex process that requires RNA polymerase II, as well as families of basal transcription factors. Binding of the factor TFIID (TBP) to the TATA box is believed to be the first step in the formation of a multiprotein complex containing several additional factors, including TFIIA, TFIIB, TFIIE, TFIIIF and TFIIH. TFIIH (or BTF2) is a multisubunit transcription/DNA repair factor that possesses several enzymatic activities. The core of TFIIH is composed of 5 subunits, designated p89 (XPB or ERCC3), p62, p52, p44 and p34. Additional subunits of the TFIIH complex are p80 (XPD or ERCC2) and the ternary kinase complex composed of Cdk7, cyclin H and MAT1. Both p89 and p80 have ATP-dependent helicase activity. The p62, p52 and p44 subunits have been shown to be involved in nucleotide excision repair.

## REFERENCES

- Conaway, R.C. and Conaway, J.W. 1989. An RNA polymerase II transcription factor has an associated DNA-dependent ATPase (dATPase) activity strongly stimulated by the TATA region of promoters. *Proc. Natl. Acad. Sci. USA* 86: 7356-7360.
- Weber, C.A., et al. 1990. ERCC2: cDNA cloning and molecular characterization of a human nucleotide excision repair gene with high homology to yeast RAD3. *EMBO J.* 9: 1437-1447.
- Weeda, G., et al. 1990. A presumed DNA helicase encoded by ERCC-3 is involved in the human repair disorders xeroderma pigmentosum and Cockayne's syndrome. *Cell* 62: 777-791.
- Fischer, L., et al. 1991. Cloning of the 62-kilodalton component of basic transcription factor BTF-2. *Science* 257: 1392-1395.
- Gerard, M., et al. 1991. Purification and interaction properties of the human polymerase B (II) general transcription factor BTF2. *J. Biol. Chem.* 266: 20940-20945.
- Flores, O., et al. 1992. Factors involved in specific transcription by mammalian RNA polymerase II. *J. Biol. Chem.* 267: 2786-2793.

## CHROMOSOMAL LOCATION

Genetic locus: *Erc2* (mouse) mapping to 7 A3.

## SOURCE

TFIIH p80 (19) is a mouse monoclonal antibody raised against amino acids 63-252 of TFIIH p80 of mouse origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-136376 X, 200 µg/0.1 ml.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

TFIIH p80 (19) is recommended for detection of TFIIH p80 of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for TFIIH p80 siRNA (m): sc-36654, TFIIH p80 shRNA Plasmid (m): sc-36654-SH and TFIIH p80 shRNA (m) Lentiviral Particles: sc-36654-V.

TFIIH p80 (19) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of TFIIH p80: 80 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210 or F9 cell lysate: sc-2245.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



TFIIH p80 (19): sc-136376. Western blot analysis of TFIIH p80 expression in F9 whole cell lysate.



TFIIH p80 (19): sc-136376. Western blot analysis of TFIIH p80 expression in NIH/3T3 whole cell lysate.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.