

CapZ- α (7): sc-136439

BACKGROUND

The F-Actin family of capping proteins includes CapZ- α 1, CapZ- α 2, CapZ- α 3 and CapZ- β 3, all of which function in a calcium-dependent manner and bind to the fast growing barbed end of Actin filaments, thereby blocking protein exchange at these ends. The F-Actin capping protein complex is a heterodimer consisting of α and β subunits that caps the barbed ends of Actin filaments and nucleates the polymerization of Actin monomers, yet does not sever Actin filaments. CapZ- α 1, also known as F-Actin-capping protein subunit α -1, is a 286 amino acid subunit of the heterodimer that forms the F-Actin capping protein complex. CapZ- α 1 also has been shown to bind S-100 β chain, a signaling molecule involved in the calcium-sensitive assembly of intermediate filaments that has been linked to Alzheimer's disease.

REFERENCES

- Casella, J.F. and Torres, M.A. 1994. Interaction of CapZ with Actin. The NH₂-terminal domains of the α 1 and β subunits are not required for Actin capping, and α 1 β and α 2 β heterodimers bind differentially to Actin. *J. Biol. Chem.* 269: 6992-6998.
- Hart, M.C., et al. 1997. Vertebrates have conserved capping protein α isoforms with specific expression patterns. *Cell Motil. Cytoskeleton* 38: 120-132.
- Inman, K.G., et al. 2002. Solution NMR structure of S100B bound to the high-affinity target peptide TRTK-12. *J. Mol. Biol.* 324: 1003-1014.
- Wear, M.A., et al. 2003. How capping protein binds the barbed end of the Actin filament. *Curr. Biol.* 13: 1531-1537.
- Hutchings, N.J., et al. 2003. Linking the T cell surface protein CD2 to the Actin-capping protein CAPZ via CMS and CIN85. *J. Biol. Chem.* 278: 22396-22403.
- Huang, S., et al. 2003. *Arabidopsis* capping protein (AtCP) is a heterodimer that regulates assembly at the barbed ends of Actin filaments. *J. Biol. Chem.* 278: 44832-44842.
- Pyle, W.G. 2004. Searching for the missing link: a role for the Actin capping protein in heart failure. *Can. J. Cardiol.* 20: 1429-1432.
- Katoh, M. 2005. WNT2B: comparative integromics and clinical applications. *Int. J. Mol. Med.* 16: 1103-1108.
- Cooper, J.A. and Sept, D. 2008. New insights into mechanism and regulation of Actin capping protein. *Int. Rev. Cell Mol. Biol.* 267: 183-206.

CHROMOSOMAL LOCATION

Genetic locus: CAPZA1 (human) mapping to 1p13.2, CAPZA2 (human) mapping to 7q31.2; Capza1 (mouse) mapping to 3 F2.2, Capza2 (mouse) mapping to 6 A2.

SOURCE

CapZ- α (7) is a mouse monoclonal antibody raised against amino acids 1-108 of CapZ- α of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CapZ- α (7) is recommended for detection of CapZ- α 1 and CapZ- α 2 of mouse, rat, human and canine origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).


Molecular Weight of CapZ- α : 36 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, CCRF-CEM cell lysate: sc-2225 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



CapZ- α (7): sc-136439. Western blot analysis of CapZ- α expression in Jurkat (A), CCRF-CEM (B), HeLa (C) and BYDP (D) whole cell lysates and rat thymus tissue extract (E).



CapZ- α (7): sc-136439. Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic staining of subset of myocytes.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures. Not for resale.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.