## SANTA CRUZ BIOTECHNOLOGY, INC.

# A1Up (T-12): sc-136560



BACKGROUND

A1Up (Ataxin-1 ubiquitin-like-interacting protein), also known as UBQLN4 (ubiquilin 4), C1orf6 or UBIN, is a 601 amino acid protein that localizes to both the cytoplasm and the nucleus and is thought to associate with the endoplasmic reticulum (ER). Expressed at high levels in kidney, pancreas, heart, brain and skeletal muscle and at lower levels in liver, lung and placenta, A1Up functions as a homodimer that binds to signal sequences on proteins that are targeted to the ER. Additionally, A1Up is thought to link Ataxin-1 with ubiquitin/proteasome pathways, possibly assisting in the Ataxin-1-associated formation of multimeric protein complexes within the nucleus. A1Up contains one ubiquitin-like domain and one UBA domain and may be phosphorylated in response to DNA damage.

### REFERENCES

- Fogli, A., Giglio, S., Arrigo, G., Lo Nigro, C., Zollo, M., Viggiano, L., Rocchi, M., Archidiacono, N., Zuffardi, O. and Carrozzo, R. 1999. Identification of two paralogous regions mapping to the short and long arms of human chromosome 2 comprising LIS1 pseudogenes. Cytogenet. Cell Genet. 86: 225-232.
- Davidson, J.D., Riley, B., Burright, E.N., Duvick, L.A., Zoghbi, H.Y. and Orr, H.T. 2000. Identification and characterization of an Ataxin-1-interacting protein: A1Up, a ubiquitin-like nuclear protein. Hum. Mol. Genet. 9: 2305-2312.
- Matsuda, M., Koide, T., Yorihuzi, T., Hosokawa, N. and Nagata, K. 2001. Molecular cloning of a novel ubiquitin-like protein, UBIN, that binds to ER targeting signal sequences. Biochem. Biophys. Res. Commun. 280: 535-540.
- 4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605440. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- Riley, B.E., Xu, Y., Zoghbi, H.Y. and Orr, H.T. 2004. The effects of the polyglutamine repeat protein Ataxin-1 on the UbL-UBA protein A1Up. J. Biol. Chem. 279: 42290-42301.

## CHROMOSOMAL LOCATION

Genetic locus: UBQLN4 (human) mapping to 1q22; UbqIn4 (mouse) mapping to 3 F1.

## SOURCE

A1Up (T-12) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the N-terminus of A1Up of human origin.

### PRODUCT

Each vial contains 100  $\mu g$  IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-136560 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

A1Up (T-12) is recommended for detection of A1Up of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:50-1:500), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:25, dilution range 1:25-1:250) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for A1Up siRNA (h): sc-78764, A1Up siRNA (m): sc-140614, A1Up shRNA Plasmid (h): sc-78764-SH, A1Up shRNA Plasmid (m): sc-140614-SH, A1Up shRNA (h) Lentiviral Particles: sc-78764-V and A1Up shRNA (m) Lentiviral Particles: sc-140614-V.


Molecular Weight of A1Up: 64-75 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, K-562 nuclear extract: sc-2130 or IMR-32 nuclear extract: sc-2148.

#### **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

## DATA



A1Up (T-12): sc-136560. Western blot analysis of A1Up expression in HeLa (A), K-562 (B), IMR-32 (C) and SH-SY-5Y (D) nuclear extracts.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### MONOS Satisfation Guaranteed Try A1Up (A333): sc-136145 or A1Up (H-2): sc-515250, our highly recommended monoclonal alternatives to A1Up (T-12).