

AURA1 (D-12): sc-136572

BACKGROUND

AURA1 (augmented in rheumatoid arthritis 1), also known as oleoyl-ACP hydrolase (OLAH), thioesterase II, S-acyl fatty acid synthase thioesterase, medium chain (SAST) or thioesterase domain containing 1 (THEDC1), is a 265 amino acid protein belonging to the thioesterase family. A component of the fatty acid synthetase complex, AURA1 catalyzes the formation of oleoyl to oleate. AURA1 expression has been found to be augmented in bone marrow-derived mononuclear cells (BMDC) of patients with rheumatoid arthritis (RA) and is encoded by a gene that maps to human chromosome 10p13, a region associated with late-onset Alzheimer disease. Making up 4.5% of the human genome, chromosome 10 encodes roughly 800 genes including PTEN, a tumor suppressor gene that has been linked to the development of Cowden syndrome. The chromosome 10 encoded gene ERCC6 is important for DNA repair and is linked to Cockayne syndrome which is characterized by extreme photosensitivity and premature aging.

REFERENCES

1. Troelstra, C., Landsvater, R.M., Wiegant, J., van der Ploeg, M., Viel, G., Buys, C.H. and Hoeijmakers, J.H. 1992. Localization of the nucleotide excision repair gene ERCC6 to human chromosome 10q11-q21. *Genomics* 12: 745-749.
2. Grupe, A., Li, Y., Rowland, C., Nowotny, P., Hinrichs, A.L., Smemo, S., Kauwe, J.S., Maxwell, T.J., Cherny, S., Doil, L., Tacey, K., van Luchene, R., Myers, A., Wavrant-De Vrièze, F., Kaleem, M., Hollingworth, P., et al 2006. A scan of chromosome 10 identifies a novel locus showing strong association with late-onset Alzheimer disease. *Am. J. Hum. Genet.* 78: 78-88.
3. Nakamura, N., Shimaoka, Y., Tougan, T., Onda, H., Okuzaki, D., Zhao, H., Fujimori, A., Yabuta, N., Nagamori, I., Tanigawa, A., Sato, J., Oda, T., Hayashida, K., Suzuki, R., Yukioka, M., Nojima, H. and Ochi, T. 2006. Isolation and expression profiling of genes upregulated in bone marrow-derived mononuclear cells of rheumatoid arthritis patients. *DNA Res.* 13: 169-183.
4. Teresi, R.E., Zbuk, K.M., Pezzolesi, M.G., Waite, K.A. and Eng, C. 2007. Cowden syndrome-affected patients with PTEN promoter mutations demonstrate abnormal protein translation. *Am. J. Hum. Genet.* 81: 756-767.
5. Yin, Y. and Shen, W.H. 2008. PTEN: a new guardian of the genome. *Oncogene* 27: 5443-5453.
6. Laugel, V., Dalloz, C., Durand, M., Sauvanaud, F., Kristensen, U., Vincent, M.C., Pasquier, L., Odent, S., Cormier-Daire, V., Gener, B., Tobias, E.S., Tolmie, J.L., Martin-Coignard, D., Drouin-Garraud, V., Heron, D., et al. 2010. Mutation update for the CSB/ERCC6 and CSA/ERCC8 genes involved in Cockayne syndrome. *Hum. Mutat.* 31: 113-126.

CHROMOSOMAL LOCATION

Genetic locus: OLAH (human) mapping to 10p13.

SOURCE

AURA1 (D-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of AURA1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-136572 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

AURA1 (D-12) is recommended for detection of AURA1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


AURA1 (D-12) is also recommended for detection of AURA1 in additional species, including equine and bovine.

Suitable for use as control antibody for AURA1 siRNA (h): sc-90654, AURA1 shRNA Plasmid (h): sc-90654-SH and AURA1 shRNA (h) Lentiviral Particles: sc-90654-V.

Molecular Weight of AURA1: 30 kDa.

Positive Controls: AURA1 (h): 293T Lysate: sc-116094.

DATA



AURA1 (D-12): sc-136572. Western blot analysis of AURA1 expression in non-transfected: sc-117752 (A) and human AURA1 transfected: sc-116094 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.