

CLUAP1 (T-14): sc-136624

BACKGROUND

CLUAP1 (Clusterin associated protein 1) is a 413 amino acid nuclear protein that exists as 2 alternatively spliced isoforms that interact with Clusterin. CLUAP1 is suggested to play a role in apoptosis and cell proliferation, and is expressed in testis, thracea and thyroid, with low levels found in adrenal gland and spinal cord. The gene encoding CLUAP1 maps to human chromosome 16, which encodes over 900 genes and comprises nearly 3% of the human genome. The GAN gene is located on chromosome 16 and, with mutation, may lead to giant axonal neuropathy, a nervous system disorder characterized by increasing malfunction with growth. The rare disorder Rubinstein-Taybi syndrome is also associated with chromosome 16, as is Crohn's disease, which is a gastrointestinal inflammatory condition.

REFERENCES

- Baraitser, M., et al. 1983. The Rubinstein-Taybi syndrome: occurrence in two sets of identical twins. *Clin. Genet.* 23: 318-320.
- Breuning, M.H., et al. 1993. Rubinstein-Taybi syndrome caused by submicroscopic deletions within 16p13.3. *Am. J. Hum. Genet.* 52: 249-254.
- Bomont, P., et al. 2000. The gene encoding gigaxonin, a new member of the cytoskeletal BTB/kelch repeat family, is mutated in giant axonal neuropathy. *Nat. Genet.* 26: 370-374.
- Kuhlenbäumer, G., et al. 2002. Giant axonal neuropathy (GAN): case report and two novel mutations in the gigaxonin gene. *Neurology* 58: 1273-1276.
- Cho, J.H. 2004. Advances in the genetics of inflammatory bowel disease. *Curr. Gastroenterol. Rep.* 6: 467-473.
- Mathew, C.G., et al. 2004. Genetics of inflammatory bowel disease: progress and prospects. *Hum. Mol. Genet.* 13: R161-R168.

CHROMOSOMAL LOCATION

Genetic locus: CLUAP1 (human) mapping to 16p13.3; Cluap1 (mouse) mapping to 16 A1.

SOURCE

CLUAP1 (T-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CLUAP1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-136624 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CLUAP1 (T-14) is recommended for detection of CLUAP1 isoforms 1 and 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CLUAP1 (T-14) is also recommended for detection of CLUAP1 isoforms 1 and 2 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for CLUAP1 siRNA (h): sc-93379, CLUAP1 siRNA (m): sc-142406, CLUAP1 shRNA Plasmid (h): sc-93379-SH, CLUAP1 shRNA Plasmid (m): sc-142406-SH, CLUAP1 shRNA (h) Lentiviral Particles: sc-93379-V and CLUAP1 shRNA (m) Lentiviral Particles: sc-142406-V.


Molecular Weight of CLUAP1: 48 kDa.

Positive Controls: CLUAP1 (h): 293T Lysate: sc-113264 or CLUAP1 (m): 293T Lysate: sc-119318.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CLUAP1 (T-14): sc-136624. Western blot analysis of CLUAP1 expression in non-transfected: sc-117752 (A) and mouse CLUAP1 transfected: sc-119318 (B) 293T whole cell lysates.



CLUAP1 (T-14): sc-136624. Western blot analysis of CLUAP1 expression in non-transfected: sc-117752 (A) and human CLUAP1 transfected: sc-113264 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.