

Clb3 (D-10): sc-136984

BACKGROUND

Cell cycle progression is controlled at a point late in G₁ designated Start. Passage through Start requires the activity of the cyclin-dependent protein kinase Cdc28. Transition from G₁/S phase requires the association of Cdc28 with members of the G₁ cyclin family, including Cln1, Cln2 and Cln3 (also designated Daf1 or Whi1). The G₂/M phase requires the M phase cyclins, Clb1 (also designated Scb1) and Clb2, as well as the G₂ cyclins, Clb3 and Clb4. The S phase cyclins Clb5 and Clb6 coordinate DNA replication with cytokinesis. Expression of the cyclins is controlled by Ubc9 and Cdc34 (also designated Udc3 or Dna6) via ubiquitin-mediated proteolysis.

REFERENCES

1. Nasmyth, K. 1993. Control of the yeast cell cycle by the Cdc28 protein kinase. *Curr. Opin. Cell Biol.* 5: 166-179.
2. Sherlock, G. and Rosamond, J. 1993. Starting to cycle: G₁ controls regulating cell division in budding yeast. *J. Gen. Microbiol.* 139: 2531-2541.
3. Amon, A., Tyers, M., Futcher, B. and Nasmyth, K. 1993. Mechanisms that help the yeast cell cycle clock tick: G₂ cyclins transcriptionally activate G₂ cyclins and repress G₁ cyclins. *Cell* 74: 993-1007.
4. Basco, R.D., Segal, M.D. and Reed, S.I. 1995. Negative regulation of G₁ and G₂ by S-phase cyclins of *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 15: 5030-5042.
5. Seufert, W., Futcher, B. and Jentsch, S. 1995. Role of a ubiquitin-conjugating enzyme in degradation of S- and M- phase cyclins. *Nature* 373: 78-81.
6. Prendergast, J.A., Ptak, C., Arnason, T.G. and Ellison, M.J. 1995. Increased ubiquitin expression suppresses the cell cycle defect associated with the yeast ubiquitin conjugating enzyme, CDC34 (UCB3). Evidence for a noncovalent interaction between CDC34 and ubiquitin. *J. Biol. Chem.* 270: 9347-9352.
7. Levine, K., Huang, K. and Cross, F.R. 1996. *Saccharomyces cerevisiae* G₁ cyclins differ in their intrinsic functional specificities. *Mol. Cell. Biol.* 16: 6794-6803.
8. Blondel, M. and Mann, C. 1996. G₂ cyclins are required for the degradation of G₁ cyclins in yeast. *Nature* 384: 279-282.

SOURCE

Clb3 (D-10) is a mouse monoclonal antibody raised against amino acids 1-427 mapping near the N-terminus of Clb3 of *Saccharomyces cerevisiae* origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS


Clb3 (D-10) is recommended for detection of Clb3 of *Saccharomyces cerevisiae* origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Clb3: 51/70 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Clb3 (D-10): sc-136984. Western blot analysis of yeast recombinant Clb3 fusion protein.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.