

# SPA-1 (C-1): sc-137043

## BACKGROUND

The SPA-1 (signal-induced proliferation-associated gene-1) protein is a principal RAP1 GTPase-activating protein in the hematopoietic progenitors and peripheral T cells. The SPA-1 gene is normally expressed in fetal and adult lymphohematopoietic tissues. Various types of mitogenic stimulation increase SPA-1 mRNA expression in normal lymphocytes. SPA-1 disrupts LFA-1-ICAM1-mediated adhesive interactions and subsequent T cell-receptor triggering and IL-2 production, possibly through inhibition of RAP1. Mice that are deficient for the SPA-1 gene develop age-dependent progression of T cell immunodeficiency followed by a spectrum of late onset myeloproliferative disorders, mimicking human chronic myeloid leukemia. SPA-1 also directly binds to AQP2 and plays a role in regulating AQP2 trafficking to the apical membrane.

## REFERENCES

1. Hattori, M., et al. 1995. Molecular cloning of a novel mitogen-inducible nuclear protein with a Ran GTPase-activating domain that affects cell cycle progression. *Mol. Cell. Biol.* 15: 552-560.
2. Katagiri, K., et al. 2002. RAP1 functions as a key regulator of T cell and antigen-presenting cell interactions and modulates T cell responses. *Mol. Cell. Biol.* 22: 1001-1015.
3. Ishida, D., et al. 2003. Antigen-driven T cell anergy and defective memory T cell response via deregulated RAP1 activation in SPA-1-deficient mice. *Proc. Natl. Acad. Sci. USA* 100: 10919-10924.
4. Harazaki, M., et al. 2004. Specific recruitment of SPA-1 to the immunological synapse: involvement of Actin-bundling protein actinin. *Immunol. Lett.* 92: 221-226.
5. Noda, Y., et al. 2004. Aquaporin-2 trafficking is regulated by PDZ-domain containing protein SPA-1. *FEBS Lett.* 568: 139-145.
6. Kometani, K., et al. 2004. RAP1 and SPA-1 in hematologic malignancy. *Trends Mol. Med.* 10: 401-408.
7. Noda, Y., et al. 2004. Molecular mechanisms and drug development in aquaporin water channel diseases: molecular mechanism of water channel aquaporin-2 trafficking. *J. Pharmacol. Sci.* 96: 249-254.
8. Noda, Y., et al. 2005. Trafficking mechanism of water channel aquaporin-2. *Biol. Cell* 97: 885-892.
9. Noda, Y., et al. 2005. Identification of a multiprotein "motor" complex binding to water channel aquaporin-2. *Biochem. Biophys. Res. Commun.* 330: 1041-1047.

## CHROMOSOMAL LOCATION

Genetic locus: SIPA1 (human) mapping to 11q13.1.

## SOURCE

SPA-1 (C-1) is a mouse monoclonal antibody raised against amino acids 691-990 mapping near the C-terminus of SPA-1 of human origin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

SPA-1 (C-1) is recommended for detection of SPA-1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SPA-1 siRNA (h): sc-45418, SPA-1 shRNA Plasmid (h): sc-45418-SH and SPA-1 shRNA (h) Lentiviral Particles: sc-45418-V.

Molecular Weight of SPA-1: 130 kDa.


Positive Controls: SPA-1 (h): 293T Lysate: sc-111824, CCRF-CEM nuclear extract: sc-2146 or Ramos cell lysate: sc-2216.

## RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.
- 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



SPA-1 (C-1): sc-137043. Western blot analysis of SPA-1 expression in non-transfected: sc-117752 (A) and human SPA-1 transfected: sc-111824 (B) 293T whole cell lysates.



SPA-1 (C-1): sc-137043. Immunoperoxidase staining of formalin fixed, paraffin-embedded human rectum tissue showing membrane and cytoplasmic staining of glandular cells.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.