

UDP-GlcDH (B-1): sc-137057

BACKGROUND

UDP-GlcDH (also called UDP-glucose 6-dehydrogenase, UGDH or UDPGDH) is a member of the UDP-glucose/GDP-mannose dehydrogenase family. UDP-GlcDH converts UDP-glucose to UDP-glucuronic acid, which is a crucial component in the biosynthesis of the glycosaminoglycans, hyaluronan, heparan sulfate and chondroitin sulfate. Found as common components of the extracellular matrix, these glycosaminoglycans are significant in signal transduction, cell migration, cancer growth and cancer metastasis. UDP-glucuronic acid (UDP-GlcA) is needed in the liver for the excretion of toxic compounds. UDP-GlcDH is an ubiquitously expressed protein most abundant in the liver. The protein structure of UDP-GlcDH was first analyzed in cow liver and found to be a homo-hexamer. This structure is well conserved between species and phyla with an overall 97% sequence identity shared between different species of mammals. Research indicates that UDP-GlcDH expression is upregulated by TGF β and downregulated by hypoxia.

CHROMOSOMAL LOCATION

Genetic locus: UGDH (human) mapping to 4p14; Ugdh (mouse) mapping to 5 C3.1.

SOURCE

UDP-GlcDH (B-1) is a mouse monoclonal antibody raised against amino acids 195-494 mapping at the C-terminus of UDP-GlcDH of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

UDP-GlcDH (B-1) is recommended for detection of UDP-GlcDH of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for UDP-GlcDH siRNA (h): sc-44709, UDP-GlcDH siRNA (m): sc-44710, UDP-GlcDH shRNA Plasmid (h): sc-44709-SH, UDP-GlcDH shRNA Plasmid (m): sc-44710-SH, UDP-GlcDH shRNA (h) Lentiviral Particles: sc-44709-V and UDP-GlcDH shRNA (m) Lentiviral Particles: sc-44710-V.


Molecular Weight of UDP-GlcDH: 57 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210, HeLa whole cell lysate: sc-2200 or C3H/10T1/2 cell lysate: sc-3801.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



UDP-GlcDH (B-1): sc-137057. Western blot analysis of UDP-GlcDH expression in HeLa (A), NIH/3T3 (B) and C3H/10T1/2 (C) whole cell lysates and rat liver tissue extract (D).



UDP-GlcDH (B-1): sc-137057. Immunoperoxidase staining of formalin fixed, paraffin-embedded human colon tissue showing nuclear and cytoplasmic staining of glandular cells.

SELECT PRODUCT CITATIONS

- Mallery, S.R., et al. 2014. Topical application of a mucoadhesive freeze-dried black raspberry gel induces clinical and histologic regression and reduces loss of heterozygosity events in premalignant oral intraepithelial lesions: results from a multicentered, placebo-controlled clinical trial. Clin. Cancer Res. 20: 1910-1924.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.