

# ERI-1 (C-8): sc-137099

## BACKGROUND

Helicase with RNase motif, more commonly designated dicer, cleaves double-stranded RNA (dsRNA) in the RNA interference and small temporal RNA (stRNA) pathways, producing active small RNA components (siRNAs) which target the destruction of RNA and repress gene expression. Human dicer cleaves dsRNA independent of ATP. The 3'-5' exonuclease ERI-1, also known as protein 3'hExo, degrades Histone mRNA after replication and may be involved in the regulation of RNA interference. ERI-1 has a high affinity for the stem-loop structure of replication-dependent Histone pre-mRNAs. It requires the 5'-ACCCA-3' sequence present in stem-loop structure. ERI-1 and a stem-loop binding protein (SLBP) target opposite faces of a unique highly conserved stem-loop RNA scaffold towards the 3' end of Histone mRNA.

## REFERENCES

- Kennedy, S., et al. 2004. A conserved siRNA-degrading RNase negatively regulates RNA interference in *C. elegans*. *Nature* 427: 645-649.
- Timmons, L. 2004. Endogenous inhibitors of RNA interference in *Caenorhabditis elegans*. *Bioessays* 26: 715-718.
- Sobering, A.K., et al. 2004. Yeast Ras regulates the complex that catalyzes the first step in GPI-anchor biosynthesis at the ER. *Cell* 117: 637-648.
- Zhang, J. 2005. Dampening the silencing effect of RNA interference in mammals. *Biochem. J.* 390: e5-e6.
- Hong, J., et al. 2005. High doses of siRNAs induce ERI-1 and ADAR1 gene expression and reduce the efficiency of RNA interference in the mouse. *Biochem. J.* 390: 675-679.
- Wang, D., et al. 2005. Somatic misexpression of germline P granules and enhanced RNA interference in retinoblastoma pathway mutants. *Nature* 436: 593-597.
- Wilkins, C., et al. 2005. RNA interference is an antiviral defence mechanism in *Caenorhabditis elegans*. *Nature* 436: 1044-1047.

## CHROMOSOMAL LOCATION

Genetic locus: ERI1 (human) mapping to 8p23.1; Eri1 (mouse) mapping to 8 A4.

## SOURCE

ERI-1 (C-8) is a mouse monoclonal antibody raised against amino acids 161-295 mapping within an internal region of ERI-1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-137099 X, 200 µg/0.1 ml.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

ERI-1 (C-8) is recommended for detection of ERI-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ERI-1 siRNA (h): sc-45559, ERI-1 siRNA (m): sc-45560, ERI-1 shRNA Plasmid (h): sc-45559-SH, ERI-1 shRNA Plasmid (m): sc-45560-SH, ERI-1 shRNA (h) Lentiviral Particles: sc-45559-V and ERI-1 shRNA (m) Lentiviral Particles: sc-45560-V.

ERI-1 (C-8) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight (predicted) of ERI-1: 34 kDa.


Molecular Weight (observed) of ERI-1: 42 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, RAW 264.7 whole cell lysate: sc-2211 or NIH/3T3 whole cell lysate: sc-2210.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



ERI-1 (C-8): sc-137099. Western blot analysis of ERI-1 expression in Jurkat (A), K-562 (B), Hep G2 (C) and 3T3-L1 (D) whole cell lysates.



ERI-1 (C-8): sc-137099. Western blot analysis of ERI-1 expression in NIH/3T3 (A), c4 (B) and RAW 264.7 (C) whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.