# SANTA CRUZ BIOTECHNOLOGY, INC.

# CPEB (E-10): sc-137146



# BACKGROUND

The regulated translation of messenger RNA is essential for cell-cycle progression, establishment of the body plan during early development, and modulation of key activities in the central nervous system. Cytoplasmic polyadenylation, one mechanism of controlling translation, is driven by cytoplasmic polyadenylation element binding protein, CPEB. CPEB is a highly conserved, sequencespecific RNA-binding protein that binds to the cytoplasmic polyadenylation element, thereby modulating translational repression and mRNA localization. Blocking cytoplasmic polyadenylation by interfering with the CPE or CPEB prevents the translational activation and translational repression of mRNAs crucial for oocyte maturation. CPEB is synthesized during oogenesis and stockpiled in the oocyte. CPEB degradation occurs via the proteasome pathway, most likely through ubiquitin-conjugated intermediates.

# **REFERENCES**

- Stebbins-Boaz, B., et al. 1999. Maskin is a CPEB-associated factor that transiently interacts with eIF-4E. Mol. Cell 4: 1017-1027.
- Luitjens, C., et al. 2000. CPEB proteins control two key steps in spermatogenesis in *C. elegans*. Genes Dev. 14: 2596-2609.

## **CHROMOSOMAL LOCATION**

Genetic locus: CPEB1 (human) mapping to 15q25.2.

# SOURCE

CPEB (E-10) is a mouse monoclonal antibody raised against amino acids 1-300 mapping at the N-terminus of CPEB of human origin.

# PRODUCT

Each vial contains 200  $\mu$ g lgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-137146 X, 200  $\mu$ g/0.1 ml.

# APPLICATIONS

CPEB (E-10) is recommended for detection of CPEB long and short isoforms of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CPEB siRNA (h): sc-37755, CPEB shRNA Plasmid (h): sc-37755-SH and CPEB shRNA (h) Lentiviral Particles: sc-37755-V.

CPEB (E-10) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of CPEB: 63 kDa.

Positive Controls: CPEB (h2): 293T Lysate: sc-116291, HeLa whole cell lysate: sc-2200 or IMR-32 cell lysate: sc-2409.

#### **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG K BP-HRP: sc-516102 or m-IgG K BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG K BP-FITC: sc-516140 or m-IgG K BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

# DATA





CPEB (E-10): sc-137146. Western blot analysis of CPEB expression in non-transfected: sc-117752 (A) and human CPEB transfected: sc-116291 (B) 2937 whole cell lysates.

CPEB (E-10): sc-137146. Western blot analysis of CPEB expression in non-transfected: sc-117752 (**A**) and human CPEB transfected: sc-114376 (**B**) 293T whole cell lysates.

#### SELECT PRODUCT CITATIONS

- Cai, C., et al. 2010. KHDC1B is a novel CPEB binding partner specifically expressed in mouse oocytes and early embryos. Mol. Biol. Cell 21: 3137-3148.
- Yu, L., et al. 2020. Neutrophil elastase-mediated proteolysis of the tumor suppressor p200 CUX1 promotes cell proliferation and inhibits cell differentiation in APL. Life Sci. 242: 117229.

# **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

# PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.