

Smt3 (D-8): sc-137177

BACKGROUND

Ubiquitin is an abundant, highly conserved protein found in all eukaryotic cells either free or covalently attached to cellular proteins. The primary function of ubiquitin in mammalian systems is to clear abnormal, foreign, and improperly folded proteins by targeting them for proteasome degradation. In *Saccharomyces cerevisiae*, ubiquitin-like proteins include Rub1, Ula1, Uba3, Smt3, Ubc2, Ubc12 and Ubc9. Rub1 shares 53% homology with ubiquitin and requires activation via Ula1, Uba3 and Ubc12 in order to conjugate to substrates directed to different proteolytic systems. Smt3, which is similar to mammalian SUMO-1, requires Ubc9 for conjugation to other proteins. Skp1 connects cell cycle regulators to the ubiquitin proteolysis machinery. Hrt1 is an essential subunit of Skp1p-cullin-F-box (SCF) complexes, which are necessary for the degradation of various regulatory proteins. Ubc13 forms a complex with Mms2 that is involved in the error-free DNA postreplication repair (PRR) pathway.

REFERENCES

1. Ciechanover, A. 1994. The ubiquitin-proteasome proteolytic pathway. *Cell* 79: 13-21.
2. Ciechanover, A. and Schwartz, A.L. 1994. The ubiquitin-mediated proteolytic pathway: mechanisms of recognition of the proteolytic substrate and involvement in the degradation of native cellular proteins. *FASEB J.* 8: 182-191.
3. Hochstrasser, M. 1995. Ubiquitin, proteasomes and the regulation of intracellular protein degradation. *Curr. Opin. Cell Biol.* 7: 215-223.
4. Bai, C., Sen, P., Hofmann, K., Ma, L., Goebel, M., Harper, J.W. and Elledge, S.J. 1996. SKP1 connects cell cycle regulators to the ubiquitin proteolysis machinery through a novel motif, the F-box. *Cell* 86: 263-274.
5. Liakopoulos, D., Doenges, G., Matuschewski, K. and Jentsch, S. 1998. A novel protein modification pathway related to the ubiquitin system. *EMBO J.* 17: 2208-2214.
6. Schwarz, S.E., Matuschewski, K., Liakopoulos, D., Scheffner, M. and Jentsch, S. 1998. The ubiquitin-like proteins SMT3 and SUMO-1 are conjugated by the UBC9 E2 enzyme. *Proc. Natl. Acad. Sci. USA* 95: 560-564.
7. Gong, L. and Yeh, E.T. 1999. Identification of the activating and conjugating enzymes of the NEDD8 conjugation pathway. *J. Biol. Chem.* 274: 12036-12042.
8. Raboy, B., Marom, A., Dor, Y. and Kulka, R.G. 1999. Heat-induced cell cycle arrest of *Saccharomyces cerevisiae*: involvement of the RAD6/UBC2 and WSC2 genes in its reversal. *Mol. Microbiol.* 32: 729-739.
9. Blondel, M., Galan, J.M. and Peter, M. 2000. Isolation and characterization of HRT1 using a genetic screen for mutants unable to degrade Gic2p in *Saccharomyces cerevisiae*. *Genetics* 155: 1033-1044.

SOURCE

Smt3 (D-8) is a mouse monoclonal antibody raised against amino acids 15-98 of Smt3 of *Saccharomyces cerevisiae* origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS


Smt3 (D-8) is recommended for detection of Smt3 of *Saccharomyces cerevisiae* origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Smt3: 12 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Smt3 (D-8): sc-137177. Western blot analysis of yeast recombinant Smt3 fusion protein.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.