SANTA CRUZ BIOTECHNOLOGY, INC.

elF3n (C-5): sc-137214



BACKGROUND

The initiation of protein synthesis in eukaryotic cells is regulated by interactions between protein initiation factors and RNA molecules. Eukaryotic initiation factors (eIFs) are utilized in a sequence of reactions that lead to 80S ribosomal assembly and, ultimately, translation. The eukaryotic initiation factor-3 (eIF3) scaffolding structure is the largest of the eIF complexes and includes eIF3 α , eIF3 β , eIF3 γ , eIF3 δ , eIF3 ε , eIF3 γ , and eIF3 θ , all of which function to control the assembly of the 40S ribosomal subunit. Association of eIF3 proteins with the 40S ribosomal subunit stabilizes eIF2-GTP-Met-tRNAiMet complex association and mRNA binding, and promotes dissociation of 80S ribosomes into 40S and 60S subunits, thereby promoting the assembly of the pre-initiation complex. Overexpression of eIF3 proteins is common in several cancers, suggesting a role for eIF3 proteins in tumorigenesis.

CHROMOSOMAL LOCATION

Genetic locus: EIF3B (human) mapping to 7p22.3; Eif3b (mouse) mapping to 5 G2.

SOURCE

elF3 η (C-5) is a mouse monoclonal antibody raised against amino acids 131-300 mapping within an internal region of elF3 η of human origin.

PRODUCT

Each vial contains 200 μg lgG1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

eIF3 η (C-5) is available conjugated to agarose (sc-137214 AC), 500 μ g/ 0.25 ml agarose in 1 ml, for IP; to HRP (sc-137214 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-137214 PE), fluorescein (sc-137214 FITC), Alexa Fluor[®] 488 (sc-137214 AF488), Alexa Fluor[®] 546 (sc-137214 AF546), Alexa Fluor[®] 594 (sc-137214 AF594) or Alexa Fluor[®] 647 (sc-137214 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-137214 AF680) or Alexa Fluor[®] 790 (sc-137214 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

elF3 η (C-5) is recommended for detection of elF3 η of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for eIF3 η siRNA (h): sc-35280, eIF3 η siRNA (m): sc-35281, eIF3 η shRNA Plasmid (h): sc-35280-SH, eIF3 η shRNA Plasmid (m): sc-35281-SH, eIF3 η shRNA (h) Lentiviral Particles: sc-35280-V and eIF3 η shRNA (m) Lentiviral Particles: sc-35281-V.

Molecular Weight of elF3₁: 116 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, NIH/3T3 whole cell lysate: sc-2210 or KNRK whole cell lysate: sc-2214.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA





eIF3 η (C-5): sc-137214. Western blot analysis of eIF3 η expression in HISM (**A**), HeLa (**B**), NIH/3T3 (**C**), RAW 264.7 (**D**), KNRK (**E**) and L8 (**F**) whole cell lysates.

elF3η (C-5): sc-137214. Immunofluorescence staining of formalin-fixed HeLa cells showing cytoplasmic localization (**A**). Immunoperoxidase staining of formalin fixed, paraffin-embedded human skin tissue showing cyto plasmic staining of keratinocytes and fibroblasts (**B**).

SELECT PRODUCT CITATIONS

- 1. Dinh, P.X., et al. 2013. Induction of stress granule-like structures in vesicular stomatitis virus-infected cells. J. Virol. 87: 372-383.
- Frydryskova, K., et al. 2016. Distinct recruitment of human elF4E isoforms to processing bodies and stress granules. BMC Mol. Biol. 17: 21.
- Chau, K.F., et al. 2018. Downregulation of ribosome biogenesis during early forebrain development. Elife 7: e36998.
- Bouillier, C., et al. 2019. The interactome analysis of the respiratory syncytial virus protein M2-1 suggests a new role in viral mRNA metabolism post-transcription. Sci. Rep. 9: 15258.
- Hans, F., et al. 2020. Multiple distinct pathways lead to hyperubiquitylated insoluble TDP-43 protein independent of its translocation into stress granules. J. Biol. Chem. 295: 673-689.
- Kim, D.Y., et al. 2020. Angiotensin AT₁ receptor antagonism by losartan stimulates adipocyte browning via induction of apelin. J. Biol. Chem. 295: 14878-14892.
- Cesaro, T., et al. 2021. PKR activity modulation by phosphomimetic mutations of serine residues located three aminoacids upstream of double-stranded RNA binding motifs. Sci. Rep. 11: 9188.
- Cascella, R., et al. 2022. A quantitative biology approach correlates neuronal toxicity with the largest inclusions of TDP-43. Sci. Adv. 8: eabm6376.
- Maharana, S., et al. 2022. SAMHD1 controls innate immunity by regulating condensation of immunogenic self RNA. Mol. Cell 82: 3712-3728.e10.
- Wang, T., et al. 2022. Intracellular energy controls dynamics of stressinduced ribonucleoprotein granules. Nat. Commun. 13: 5584.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA