

ACSVL3 (Q-12): sc-137275

BACKGROUND

Acyl-coenzyme A synthetases (ACSSs) are a large family of related enzymes known to catalyze the fundamental initial reaction in fatty acid metabolism. The ACS family is roughly characterized based on fatty acid chain length preference amongst different members. The nomenclature in the ACS family reflects this relationship and includes short-chain ACS (ACSS), medium-chain ACS (ACSM), long-chain ACS (ACSL) and very long-chain ACS (ACSVL). ACSVL family members are capable of activating both long (LCFAs) and very long-chain fatty acids (VLCFAs). There are six members of the human ACSVL sub-family, which have been described as solute carrier family 27A (SLC27A) gene products. They represent a group of evolutionarily conserved fatty acid transport proteins (FATPs) recognized for their role in facilitating translocation of long-chain fatty acids across the plasma membrane. The family nomenclature has recently been unified with their respective acyl-CoA synthetase family designations: ACSVL1 (FATP2), ACSVL2 (FATP6), ACSVL3 (FATP3), ACSVL4 (FATP1), ACSVL5 (FATP4) and ACSVL6 (FATP5). ACSVLs have unique expression patterns and are found in major organs of fatty acid metabolism, such as adipose tissue, liver, heart and kidney. ACSVL3 is a 730 amino acid multi-pass membrane protein. Encoded by a gene that maps to human chromosome 1q21.3, ACSVL3 exists as three alternatively spliced isoforms.

REFERENCES

- Schaffer, J.E., et al. 1994. Expression cloning and characterization of a novel adipocyte long chain fatty acid transport protein. *Cell* 79: 427-436.
- Hirsch, D., et al. 1998. A family of fatty acid transporters conserved from mycobacterium to man. *Proc. Natl. Acad. Sci. USA* 95: 8625-8629.
- Abumrad, N., et al. 1999. Membrane proteins implicated in long-chain fatty acid uptake by mammalian cells: CD36, FATP and FABPm. *Biochim. Biophys. Acta* 1441: 4-13.
- Binnert, C., et al. 2000. Fatty acid transport protein-1 mRNA expression in skeletal muscle and in adipose tissue in humans. *Am. J. Physiol. Endocrinol. Metab.* 279: E1072-E1079.
- Martin, G., et al. 2000. The human fatty acid transport protein-1 (SLC27A1; FATP-1) cDNA and gene: organization, chromosomal localization, and expression. *Genomics* 66: 296-304.

CHROMOSOMAL LOCATION

Genetic locus: SLC27A3 (human) mapping to 1q21.3; Slc27a3 (mouse) mapping to 3 F1.

SOURCE

ACSVL3 (Q-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ACSVL3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-137275 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ACSVL3 (Q-12) is recommended for detection of ACSVL3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ACSVL family members.

ACSVL3 (Q-12) is also recommended for detection of ACSVL3 in additional species, including bovine.

Suitable for use as control antibody for ACSVL3 siRNA (h): sc-78881, ACSVL3 siRNA (m): sc-140838, ACSVL3 shRNA Plasmid (h): sc-78881-SH, ACSVL3 shRNA Plasmid (m): sc-140838-SH, ACSVL3 shRNA (h) Lentiviral Particles: sc-78881-V and ACSVL3 shRNA (m) Lentiviral Particles: sc-140838-V.


Molecular Weight of ACSVL3: 79 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, Hep G2 cell lysate: sc-2227 or U-251-MG whole cell lysate: sc-364176.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



ACSVL3 (Q-12): sc-137275. Western blot analysis of ACSVL3 expression in K-562 (A), U-251-MG (B) and Hep G2 (C) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.