

DPAGT1 (D-15): sc-137436

BACKGROUND

DPAGT1 (UDP-N-acetylglucosamine-dolichyl-phosphate N-acetylglucosaminophosphotransferase), also known as G1PT (GlcNAc-1-P transferase) or DPAGT2, is a 408 amino acid member of the glycosyltransferase 4 protein family. Localized to the endoplasmic reticulum membrane, DPAGT1 is involved in protein modification, specifically glycosylation. DPAGT1 catalyzes the initial step in the synthesis of dolichol-P-P-oligosaccharides. Defects in the gene that encodes DPAGT1 are the cause of congenital disorder of glycosylation type 1J (CDG1J). Congenital disorders of glycosylation (CDGs) are a family of severe inherited diseases caused by a defect in protein N-glycosylation. CDGs cause a variety of clinical features including dysmorphic features, psychomotor retardation, hypotonia, coagulation disorders and immunodeficiency.

REFERENCES

- Eckert, V., et al. 1998. Cloning and functional expression of the human GlcNAc-1-P transferase, the enzyme for the committed step of the dolichol cycle, by heterologous complementation in *Saccharomyces cerevisiae*. *Glycobiology* 8: 77-85.
- Meissner, J.D., et al. 1999. Regulation of UDP-N-acetylglucosamine: dolichyl-phosphate N-acetylglucosamine-1-phosphate transferase by retinoic acid in P19 cells. *Biochem. J.* 338: 561-568.
- Freeze, H.H. 2002. Human disorders in N-glycosylation and animal models. *Biochim. Biophys. Acta* 1573: 388-393.
- Regis, S., et al. 2002. Genomic structure of the human UDP-GlcNAc: dolichol-P GlcNAc-1-P transferase gene. *DNA Seq.* 13: 245-250.
- Wu, X., et al. 2003. Deficiency of UDP-GlcNAc:dolichol phosphate N-acetylglucosamine-1 phosphate transferase (DPAGT1) causes a novel congenital disorder of Glycosylation Type 1j. *Hum. Mutat.* 22: 144-150.
- Newell, J.W., et al. 2003. Congenital disorder of glycosylation 1c in patients of Indian origin. *Mol. Genet. Metab.* 79: 221-228.
- Nita-Lazar, M., et al. 2009. Overexpression of DPAGT1 leads to aberrant N-glycosylation of E-cadherin and cellular dis-cohesion in oral cancer. *Cancer Res.* 69: 5673-5680.
- Bretthauer, R.K. 2009. Structure, expression, and regulation of UDP-GlcNAc: dolichol phosphate GlcNAc-1-phosphate transferase (DPAGT1). *Curr. Drug Targets* 10: 477-482.
- Sengupta, P.K., et al. 2010. N-glycosylation gene DPAGT1 is a target of the Wnt/ β -catenin signaling pathway. *J. Biol. Chem.* 285: 31164-31173.

CHROMOSOMAL LOCATION

Genetic locus: DPAGT1 (human) mapping to 11q23.3; Dpagt1 (mouse) mapping to 9 A5.2.

SOURCE

DPAGT1 (D-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal cytoplasmic domain of DPAGT1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-137436 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

DPAGT1 (D-15) is recommended for detection of DPAGT1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

DPAGT1 (D-15) is also recommended for detection of DPAGT1 in additional species, including equine, bovine, porcine and avian.

Suitable for use as control antibody for DPAGT1 siRNA (h): sc-96745, DPAGT1 siRNA (m): sc-143149, DPAGT1 shRNA Plasmid (h): sc-96745-SH, DPAGT1 shRNA Plasmid (m): sc-143149-SH, DPAGT1 shRNA (h) Lentiviral Particles: sc-96745-V and DPAGT1 shRNA (m) Lentiviral Particles: sc-143149-V.

Molecular Weight of DPAGT1 isoforms: 46/34 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.