

# UBE2F (S-12): sc-137893

## BACKGROUND

Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). UBE2F (ubiquitin-conjugating enzyme E2F), also known as NCE2, is a 185 amino acid protein that belongs to the E2 family of ubiquitin conjugating enzymes. UBE2F functions to accept the ubiquitin-like protein NEDD8 from an E1 complex and to catalyze the ATP-dependent attachment of NEDD8 to other proteins, playing a role in the pathway of protein degradation. The gene encoding UBE2F maps to chromosome 2, which encodes over 1,400 genes and comprises nearly 8% of the human genome.

## CHROMOSOMAL LOCATION

Genetic locus: UBE2F (human) mapping to 2q37.3; Ube2f (mouse) mapping to 1 D.

## SOURCE

UBE2F (S-12) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the N-terminus of UBE2F of human origin.

## PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-137893 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

UBE2F (S-12) is recommended for detection of UBE2F of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:50-1:500), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:25, dilution range 1:25-1:250) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other UBE2 family members.

UBE2F (S-12) is also recommended for detection of UBE2F in additional species, including canine and avian.

Suitable for use as control antibody for UBE2F siRNA (h): sc-94988, UBE2F siRNA (m): sc-154852, UBE2F shRNA Plasmid (h): sc-94988-SH, UBE2F shRNA Plasmid (m): sc-154852-SH, UBE2F shRNA (h) Lentiviral Particles: sc-94988-V and UBE2F shRNA (m) Lentiviral Particles: sc-154852-V.


Molecular Weight of UBE2F isoforms: 24/21/18/14/11 kDa.

Positive Controls: UBE2F (h): 293 Lysate: sc-112735, UBE2F (m): 293T Lysate: sc-124413 or Jurkat whole cell lysate: sc-2204.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



UBE2F (S-12): sc-137893. Western blot analysis of UBE2F expression in non-transfected: sc-110760 (A) and human UBE2F transfected: sc-112735 (B) 293 whole cell lysates.



UBE2F (S-12): sc-137893. Western blot analysis of UBE2F expression in non-transfected: sc-117752 (A) and mouse UBE2F transfected: sc-124413 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **UBE2F (C-11): sc-398668**, our highly recommended monoclonal alternative to UBE2F (S-12).