

# HABP4 (P-13): sc-138020

## BACKGROUND

The intracellular hyaluronan-binding protein 4 (HABP4 or iHABP4), also designated Ki-1/57, is a cytoplasmic and nuclear protein that is phosphorylated on serine and threonine residues. HABP4 is expressed as a protein of 413 amino acids and is phosphorylated by phorbol 12-myristate 13-acetate (PMA)-activated PKC isoforms at Thr-354 and Thr-375. HABP4 interacts with hyaluron, the adaptor protein RACK1, the transcription factor MEF2C, and the chromatin remodeling factor CHD3, suggesting that it might be involved in the regulation of transcription. However, its actual function has yet to be elucidated. HABP4 is highly expressed in brain, heart, and kidney, and moderately expressed in skeletal muscle. Also expressed in a variety of tumor cell lines and in activated but not resting leukocytes.

## REFERENCES

1. Kobarg, J., Schnittger, S., Fonatsch, C., Lemke, H., Bowen, M.A., Buck, F. and Hansen, H.P. 1997. Characterization, mapping and partial cDNA sequence of the 57-kD intracellular Ki-1 antigen. *Exp. Clin. Immunogenet.* 14: 273-280.
2. Huang, L., Grammatikakis, N., Yoneda, M., Banerjee, S.D. and Toole, B.P. 2000. Molecular characterization of a novel intracellular hyaluronan-binding protein. *J. Biol. Chem.* 275: 29829-29839.
3. Lemos, T.A., Passos, D.O., Nery, F.C. and Kobarg, J. 2003. Characterization of a new family of proteins that interact with the C-terminal region of the chromatin-remodeling factor CHD-3. *FEBS Lett.* 533: 14-20.
4. Nery, F.C., Passos, D.O., Garcia, V.S. and Kobarg, J. 2004. Ki-1/57 interacts with RACK1 and is a substrate for the phosphorylation by phorbol 12-myristate 13-acetate-activated protein kinase C. *J. Biol. Chem.* 279: 11444-11455.
5. Kobarg, C.B., Kobarg, J., Crosara-Alberto, D.P., Theizen, T.H. and Franchini, K.G. 2005. MEF2C DNA-binding activity is inhibited through its interaction with the regulatory protein Ki-1/57. *FEBS Lett.* 579: 2615-2622.
6. Nery, F.C., Rui, E., Kuniyoshi, T.M. and Kobarg, J. 2006. Evidence for the interaction of the regulatory protein Ki-1/57 with p53 and its interacting proteins. *Biochem. Biophys. Res. Commun.* 341: 847-855.
7. Passos, D.O., Bressan, G.C., Nery, F.C. and Kobarg, J. 2006. Ki-1/57 interacts with PRMT1 and is a substrate for arginine methylation. *FEBS J.* 273: 3946-3961.
8. Bressan, G.C., Silva, J.C., Borges, J.C., Dos Passos, D.O., Ramos, C.H., Torriani, I.L. and Kobarg, J. 2008. Human regulatory protein Ki-1/57 has characteristics of an intrinsically unstructured protein. *J. Proteome Res.* 7: 4465-4474.
9. Bressan, G.C., Quaresma, A.J., Moraes, E.C., Manfiolli, A.O., Passos, D.O., Gomes, M.D. and Kobarg, J. 2009. Functional association of human Ki-1/57 with pre-mRNA splicing events. *FEBS J.* 276: 3770-3783.

## CHROMOSOMAL LOCATION

Genetic locus: HABP4 (human) mapping to 9q22.32; Habp4 (mouse) mapping to 13 B3.

## SOURCE

HABP4 (P-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of HABP4 of mouse origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-138020 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

HABP4 (P-13) is recommended for detection of HABP4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with HABP2.


Suitable for use as control antibody for HABP4 siRNA (h): sc-92886, HABP4 siRNA (m): sc-145888, HABP4 shRNA Plasmid (h): sc-92886-SH, HABP4 shRNA Plasmid (m): sc-145888-SH, HABP4 shRNA (h) Lentiviral Particles: sc-92886-V and HABP4 shRNA (m) Lentiviral Particles: sc-145888-V.

Molecular Weight (predicted) of HABP4: 46 kDa.

Molecular Weight (observed) of HABP4: 53 kDa.

Positive Controls: U-87 MG cell lysate: sc-2411.

## DATA



HABP4 (P-13): sc-138020. Western blot analysis of HABP4 expression in U-87 MG whole cell lysate.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.