

EF-CAB7 (N-13): sc-138255

BACKGROUND

The intracellular calcium-binding superfamily of proteins consists of EF-hand calcium binding domains and are often involved in the regulation of many different cellular processes. An EF-hand calcium binding domain is made up of approximately 40 amino acids and can bind to 2 intracellular calcium ions. Two types of EF-hand calcium binding motifs exist: regulatory and structural. Proteins containing the regulatory EF-hand domain induce conformational change, allowing interaction with target proteins and catalyzing enzymatic reactions, whereas structural EF-hand domain containing proteins do not undergo conformational change and may play a role in buffering intracellular calcium levels. EF-CAB7 (EF-hand calcium binding domain 7) is a 629 amino acid protein that contains 3 EF-hand domains. EF-CAB7 contains two alternatively spliced isoforms and is encoded by a gene located on human chromosome 1p31.3.

REFERENCES

1. Babitch, J.A. et al. 1987. Grasping for calcium binding sites in sodium channels with an EF hand. *J. Theor. Biol.* 127: 451-459.
2. Perret, C., et al. 1988. Evolution of the EF-hand calcium-binding protein family: evidence for exon shuffling and intron insertion. *J. Mol. Evol.* 27: 351-364.
3. Taylor, D.A., et al. 1991. Structure of a recombinant calmodulin from *Drosophila melanogaster* refined at 2.2-Å resolution. *J. Biol. Chem.* 266: 21375-21380.
4. Drake, S.K., et al. 1996. Tuning the equilibrium ion affinity and selectivity of the EF-hand calcium binding motif: substitutions at the gateway position. *Biochemistry* 35: 6697-6705.
5. Drake, S.K., et al. 1997. Molecular tuning of an EF-hand-like calcium binding loop. Contributions of the coordinating side chain at loop position 3. *J. Gen. Physiol.* 110: 173-184.
6. Atkinson, R.A., et al. 2001. Ca²⁺-independent binding of an EF-hand domain to a novel motif in the α -actinin-titin complex. *Nat. Struct. Biol.* 8: 853-857.
7. Julenius, K., et al. 2002. Coupling of ligand binding and dimerization of helix-loop-helix peptides: spectroscopic and sedimentation analyses of calbindin D9k EF-hands. *Proteins* 47: 323-333.

CHROMOSOMAL LOCATION

Genetic locus: EFCAB7 (human) mapping to 1p31.3.

SOURCE

EF-CAB7 (N-13) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping at the N-terminus of EF-CAB7 of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-138255 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

EF-CAB7 (N-13) is recommended for detection of EF-CAB7 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:50-1:500), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:25, dilution range 1:25-1:250) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other EF-CAB family members.

Suitable for use as control antibody for EF-CAB7 siRNA (h): sc-88682, EF-CAB7 shRNA Plasmid (h): sc-88682-SH and EF-CAB7 shRNA (h) Lentiviral Particles: sc-88682-V.

Molecular Weight of EF-CAB7 isoform 1: 72 kDa.


Molecular Weight of EF-CAB7 isoform 2: 69 kDa.

Positive Controls: EF-CAB7 (h): 293T Lysate: sc-112202.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



EF-CAB7 (N-13): sc-138255. Western blot analysis of EF-CAB7 expression in non-transfected: sc-117752 (A) and human EF-CAB7 transfected: sc-112202 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.