UVSSA (P-12): sc-138374



The Power to Question

BACKGROUND

UVSSA (UV-stimulated scaffold protein A), also known as KIAA1530, is a 709 amino acid chromosomal protein belonging to the UVSSA family and containing one coiled coil domain and one VHS-like region. Defects in the UVSSA gene result in UV-sensitive syndrome 3 (UVSS3), an autosomal recessive disorder characterized by cutaneous photosensitivity and slight dyspigmentation without skin carcinoma. UVSSA interacts with the machinery for transcription-coupled nucleotide-excision repair (TC-NER) which allows RNA polymerase Il-blocking lesions to be removed from the transcribed strand of active genes. UVSSA also stabilizes the ERCC6 (excision repair cross-complementation group 6) complex, and facilitates ubiquitination of the elongating form of RNA polymerase II (RNA pol IIo) stalled at DNA damage sites.

REFERENCES

- 1. Yashin, A.I., et al. 2010. Joint influence of small-effect genetic variants on human longevity. Aging 2: 612-620.
- Fei, J., et al. 2012. KIAA1530 protein is recruited by Cockayne syndrome complementation group protein A (CSA) to participate in transcriptioncoupled repair (TCR). J. Biol. Chem. 287: 35118-35126.
- 3. Nakazawa, Y., et al. 2012. Mutations in UVSSA cause UV-sensitive syndrome and impair RNA polymerase IIo processing in transcription-coupled nucleotide-excision repair. Nat. Genet. 44: 586-592.
- 4. Zhang, X., et al. 2012. Mutations in UVSSA cause UV-sensitive syndrome and destabilize ERCC6 in transcription-coupled DNA repair. Nat. Genet. 44: 593-597.
- Schwertman, P., et al. 2012. UV-sensitive syndrome protein UVSSA recruits USP7 to regulate transcription-coupled repair. Nat. Genet. 44: 598-602.

CHROMOSOMAL LOCATION

Genetic locus: UVSSA (human) mapping to 4p16.3; UVSSA (mouse) mapping to 5 $\rm B1$.

SOURCE

UVSSA (P-12) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the C-terminus of UVSSA of human origin.

PRODUCT

Each vial contains 100 μg lgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-138374 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

UVSSA (P-12) is recommended for detection of UVSSA of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:50-1:500), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:25, dilution range 1:25-1:250) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for UVSSA siRNA (h): sc-89002, UVSSA siRNA (m): sc-149456, UVSSA shRNA Plasmid (h): sc-89002-SH, UVSSA shRNA Plasmid (m): sc-149456-SH, UVSSA shRNA (h) Lentiviral Particles: sc-89002-V and UVSSA shRNA (m) Lentiviral Particles: sc-149456-V.


Molecular Weight of UVSSA isoforms 1/2: 81/30 kDa.

Positive Controls: Y79 cell lysate: sc-2240, MOLT-4 cell lysate: sc-2233 or Raji whole cell lysates: sc-364236

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



UVSSA (P-12): sc-138374. Western blot analysis of UVSSA expression in Y79 (A), MOLT-4 (B), Raji (C) whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.