

FBXO30 (P-20): sc-138936

BACKGROUND

FBXO30 (F-box only protein 30) is a 745 amino acid protein that contains one 40 amino acid F-box region, making it a member of the F-box family. FBXO30 also contains one TRAF-type zinc finger. F-box proteins are critical components of the SCF (Skp1-CUL-1-F-box protein) type E3 ubiquitin ligase complex and are involved in substrate recognition and recruitment for ubiquitination. F-box proteins are members of a large family that regulates cell cycle, immune response, signaling cascades and developmental programs by targeting proteins, such as cyclins, cyclin-dependent kinase inhibitors, κ B- α and β -catenin, for degradation by the proteasome after ubiquitination. Functioning as a component of the SCF complex, FBXO30 is thought to recognize and bind to select phosphorylated proteins, thereby promoting their ubiquitination and subsequent degradation.

REFERENCES

1. Winston, J.T., et al. 1999. The SCF β -TrCP-ubiquitin ligase complex as-sociates specifically with phosphorylated destruction motifs in κ B- α and β -catenin and stimulates κ B- α ubiquitination *in vitro*. *Genes Dev.* 13: 270-283.
2. Cenciarelli, C., et al. 1999. Identification of a family of human F-box proteins. *Curr. Biol.* 9: 1177-1179.
3. Winston, J.T., et al. 1999. A family of mammalian F-box proteins. *Curr. Biol.* 9: 1180-1182.
4. Craig, K.L., et al. 1999. The F-box: a new motif for ubiquitin dependent proteolysis in cell cycle regulation and signal transduction. *Prog. Biophys. Mol. Biol.* 72: 299-328.
5. Ilyin, G.P., et al. 2000. cDNA cloning and expression analysis of new members of the mammalian F-box protein family. *Genomics* 67: 40-47.
6. Schulman, B.A., et al. 2000. Insights into SCF ubiquitin ligases from the structure of the Skp1-Skp2 complex. *Nature* 408: 381-386.
7. Ilyin, G.P., et al. 2002. A new subfamily of structurally related human F-box proteins. *Gene* 296: 11-20.

CHROMOSOMAL LOCATION

Genetic locus: FBXO30 (human) mapping to 6q24.3.

SOURCE

FBXO30 (P-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of FBXO30 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-138936 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

FBXO30 (P-20) is recommended for detection of FBXO30 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other FBXO family members.

Suitable for use as control antibody for FBXO30 siRNA (h): sc-95455, FBXO30 shRNA Plasmid (h): sc-95455-SH and FBXO30 shRNA (h) Lentiviral Particles: sc-95455-V.


Molecular Weight of FBXO30: 82 kDa.

Positive Controls: FBXO30 (h): 293T Lysate: sc-113671.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



FBXO30 (P-20): sc-138936. Western blot analysis of FBXO30 expression in non-transfected: sc-117752 (A) and human FBXO30 transfected: sc-113671 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

MONOS
Satisfaction
Guaranteed

Try **FBXO30 (E-10): sc-514862**, our highly recommended monoclonal alternative to FBXO30 (P-20).