


# ASIC1 (N-18): sc-13909



## BACKGROUND

Degenerin/epithelial sodium channel (DEG/ENaC) superfamily members are amiloride-sensitive sodium channels that contain intracellular N- and C-terminal, two hydrophobic transmembrane regions and a cysteine-containing extracellular loop. Acid sensing ion channel ASIC1, also designated ACCN2, BNAC2 and ASIC1α, is present in brain as a 4.3-kb transcript with localization to rat dorsal root ganglia. *In situ* hybridization of rat brain suggests that ASIC1 is most abundant in the main olfactory bulb, cerebral cortex, hippocampal formation, habenula, basolateral amygdaloid nuclei and cerebellum. ASIC1 and H<sup>+</sup>-gated currents may contribute to the development of fear and anxiety. ASIC2, also designated amiloride-sensitive cation channel 1, neuronal (ACCN1), mammalian degenerin, BNAC1 (MDEG) and brain Na<sup>+</sup> channel 1, mediates the normal detection of light touch. ASIC2 mRNA is abundant in brain, specifically in neurons. ASIC2 is expressed as 2.7- and 3.7-kb transcripts in brain and spinal cord tissues. ASIC3, also designated SLNAC1 and TNAC1, mediates detection of lasting pH changes and is involved in modulating moderate- to high-intensity pain sensation. ASIC4, also designated ACCN4 and BNAC4, is abundant in pituitary gland and is also present in the inner ear.

## REFERENCES

- Waldmann, R., et al. 1997. A proton-gated cation channel involved in acid-sensing. *Nature* 386: 173-177.
- Garcia-Anoveros, J., et al. 1997. BNAC1 and BNAC2 constitute a new family of human neuronal sodium channels related to degenerins and epithelial sodium channels. *Proc. Natl. Acad. Sci. USA* 94: 1459-1464.
- Price, M.P., et al. 2000. The mammalian sodium channel BNC1 is required for normal touch sensation. *Nature* 407: 1007-1011.
- Grunder, S., et al. 2001. Acid-sensing ion channel (ASIC) 4 gene: physical mapping, genomic organisation, and evaluation as a candidate for paroxysmal dystonia. *Eur. J. Hum. Genet.* 9: 672-676.
- Chen, C.C., et al. 2002. A role for ASIC3 in the modulation of high-intensity pain stimuli. *Proc. Natl. Acad. Sci. USA* 99: 8992-8997.

## CHROMOSOMAL LOCATION

Genetic locus: ASIC1 (human) mapping to 12q13.12.

## SOURCE

ASIC1 (N-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ASIC1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-13909 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

ASIC1 (N-18) is recommended for detection of ASIC1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with ASIC-β.

Suitable for use as control antibody for ASIC1 siRNA (h): sc-42407, ASIC1 shRNA Plasmid (h): sc-42407-SH and ASIC1 shRNA (h) Lentiviral Particles: sc-42407-V.


Molecular Weight of ASIC1: 60 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409, HT-1080 whole cell lysate: sc-364183 or CCRF-CEM cell lysate: sc-2225.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



ASIC1 (N-18): sc-13909. Western blot analysis of ASIC1 expression in IMR-32 (**A**), U-87 MG (**B**), HT-1080 (**C**), CCRF-CEM (**D**) and HEK293 (**E**) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.