

NAT-8 (L-12): sc-139400

BACKGROUND

Acetyltransferases and deacetylases are protein groups most often associated with oncogenesis and cell cycle regulation. NAT-8 (N-acetyltransferase 8), also known as GLA, CML1 (camello-like protein 1) or TSC510, is a 227 amino acid multi-pass membrane protein that plays a role in gastrulation regulation. A member of the camello family, NAT-8 contains one N-acetyltransferase domain and is expressed in kidney and liver. The gene encoding NAT-8 maps to human chromosome 2, which consists of 237 million bases, encodes over 1,400 genes and makes up approximately 8% of the human genome. A number of genetic diseases are linked to genes on chromosome 2 including Harlequin ichthyosis, sitosterolemia and Alström syndrome.

REFERENCES

- Patel, S.B., et al. 1998. Mapping a gene involved in regulating dietary cholesterol absorption. The sitosterolemia locus is found at chromosome 2p21. *J. Clin. Invest.* 102: 1041-1044.
- Ozaki, K., et al. 1998. Isolation and mapping of a novel human kidney- and liver-specific gene homologous to the bacterial acetyltransferases. *J. Hum. Genet.* 43: 255-258.
- Zumsteg, U., et al. 2000. Alstrom syndrome: confirmation of linkage to chromosome 2p12-13 and phenotypic heterogeneity in three affected sibs. *J. Med. Genet.* 37: E8.
- Popsueva, A.E., et al. 2001. Overexpression of camello, a member of a novel protein family, reduces blastomere adhesion and inhibits gastrulation in *Xenopus laevis*. *Dev. Biol.* 234: 483-496.
- Kelsell, D.P., et al. 2005. Mutations in ABCA12 underlie the severe congenital skin disease harlequin ichthyosis. *Am. J. Hum. Genet.* 76: 794-803.
- Juhanson, P., et al. 2008. N-acetyltransferase 8, a positional candidate for blood pressure and renal regulation: resequencing, association and in silico study. *BMC Med. Genet.* 9: 25.
- Ariyannur, P.S., et al. 2010. Methamphetamine-induced neuronal protein NAT8L is the NAA biosynthetic enzyme: implications for specialized acetyl coenzyme A metabolism in the CNS. *Brain Res.* 1335: 1-13.
- Veiga-da-Cunha, M., et al. 2010. Molecular identification of NAT8 as the enzyme that acetylates cysteine S-conjugates to mercapturic acids. *J. Biol. Chem.* 285: 18888-18898.
- Suhre, K., et al. 2011. Human metabolic individuality in biomedical and pharmaceutical research. *Nature* 477: 54-60.

CHROMOSOMAL LOCATION

Genetic locus: NAT8 (human) mapping to 2p13.1.

SOURCE

NAT-8 (L-12) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of NAT-8 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-139400 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

NAT-8 (L-12) is recommended for detection of NAT-8 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:50-1:500), immunofluorescence (starting dilution 1:25, dilution range 1:25-1:250) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); may cross-react with NAT-8B.

Suitable for use as control antibody for NAT-8 siRNA (h): sc-94277, NAT-8 shRNA Plasmid (h): sc-94277-SH and NAT-8 shRNA (h) Lentiviral Particles: sc-94277-V.

Molecular Weight of NAT-8: 26 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.