

VMO1 (S-18): sc-139446

BACKGROUND

VMO1 [vitelline membrane outer layer 1 homolog (chicken)] is a 202 amino acid secreted protein that belongs to the VMO1 family. VMO1 is encoded by a gene located on human chromosome 17p13.2. Chromosome 17 makes up over 2.5% of the human genome, with about 81 million bases encoding over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, though it is specifically recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes. Chromosome 17 is also linked to neurofibromatosis, a condition characterized by neural and epidermal lesions, and dysregulated Schwann cell growth. Alexander disease, Birt-Hogg-Dube syndrome and Canavan disease are also associated with chromosome 17.

REFERENCES

1. Welsch, M.J., et al. 2005. Birt-Hogg-Dube Syndrome. *Int. J. Dermatol.* 44: 668-673.
2. Nusbaum, R., et al. 2006-2007. Susceptibility to breast cancer: hereditary syndromes and low penetrance genes. *Breast Dis.* 27: 21-50.
3. Al-Dirbashi, O.Y., et al. 2007. Quantification of N-acetylaspartic acid in urine by LC-MS/MS for the diagnosis of Canavan disease. *J. Inher. Metab. Dis.* 30: 612.
4. Dann, R.B., et al. 2007. Strategies for ovarian cancer prevention. *Obstet. Gynecol. Clin. North Am.* 34: 667-686.

CHROMOSOMAL LOCATION

Genetic locus: VMO1 (human) mapping to 17p13.2; Vmo1 (mouse) mapping to 11 B3.

SOURCE

VMO1 (S-18) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of VMO1 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-139446 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

VMO1 (S-18) is recommended for detection of VMO1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:50-1:500), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:25, dilution range 1:25-1:250), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:25, dilution range 1:25-1:250) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

VMO1 (S-18) is also recommended for detection of VMO1 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for VMO1 siRNA (h): sc-93959, VMO1 siRNA (m): sc-155216, VMO1 shRNA Plasmid (h): sc-93959-SH, VMO1 shRNA Plasmid (m): sc-155216-SH, VMO1 shRNA (h) Lentiviral Particles: sc-93959-V and VMO1 shRNA (m) Lentiviral Particles: sc-155216-V.


Molecular Weight of VMO1: 22 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or VMO1 (m): 293T Lysate: sc-179738.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



VMO1 (S-18): sc-139446. Western blot analysis of VMO1 expression in non-transfected 293T: sc-117752 (A), mouse VMO1 transfected 293T: sc-179738 (B) and HeLa (C) whole cell lysates.



VMO1 (S-18): sc-139446. Immunoperoxidase staining of formalin fixed, paraffin-embedded human epididymis tissue showing cytoplasmic and membrane staining of glandular cells.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.