

α -sarcoglycan (C-15): sc-14174

BACKGROUND

The sarcoglycan transmembrane proteins are members of the dystrophin complex. Sarcoglycans cluster together to form a complex, which is localized in the cell membrane of skeletal, cardiac, and smooth muscle fibers. Four sarcoglycan subunit proteins, designated α -, β -, γ - and δ -sarcoglycan, form a complex on the skeletal muscle cell surface membrane. A genetic defect in any one of these proteins causes the loss or marked decrease of the whole sarcoglycan complex, which is observed in the autosomal recessive muscular dystrophy, sarcoglycanopathy. In smooth muscle, β - and δ -sarcoglycans are associated with ϵ -sarcoglycan, a glycoprotein homologous to α -sarcoglycan. Additionally, a complete deficiency in δ -sarcoglycan is the cause of the Syrian hamster BIO.14 cardiomyopathy.

REFERENCES

1. Barresi, R., et al. 2000. Expression of γ -sarcoglycan in smooth muscle and its interaction with the smooth muscle sarcoglycan-sarcospan complex. *J. Biol. Chem.* 275: 38554-38560.
2. Hack, A.A., et al. 2000. Differential requirement for individual sarcoglycans and dystrophin in the assembly and function of the dystrophin-glycoprotein complex. *J. Cell Sci.* 113: 2535-2544.
3. Ueda, H., et al. 2001. δ - and γ -sarcoglycan localization in the sarcoplasmic reticulum of skeletal muscle. *J. Histochem. Cytochem.* 49: 529-538.
4. Wakabayashi-Takai, E., et al. 2001. Identification of myogenesis-dependent transcriptional enhancers in promoter region of mouse γ -sarcoglycan gene. *Eur. J. Biochem.* 268: 948-957.
5. Politano, L., et al. 2001. Evaluation of cardiac and respiratory involvement in sarcoglycanopathies. *Neuromuscul. Disord.* 11: 178-185.
6. Enigk, R.E., et al. 2001. Cellular and molecular properties of α -dystrobrevin in skeletal muscle. *Front. Biosci.* 6: 53-64.

CHROMOSOMAL LOCATION

Genetic locus: SGCA (human) mapping to 17q21.33; Sgca (mouse) mapping to 11 D.

SOURCE

α -sarcoglycan (C-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of α -sarcoglycan of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-14174 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

α -sarcoglycan (C-15) is recommended for detection of α -sarcoglycan of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

α -sarcoglycan (C-15) is also recommended for detection of α -sarcoglycan in additional species, including canine.

Suitable for use as control antibody for α -sarcoglycan siRNA (h): sc-43416, α -sarcoglycan siRNA (m): sc-43417, α -sarcoglycan shRNA Plasmid (h): sc-43416-SH, α -sarcoglycan shRNA Plasmid (m): sc-43417-SH, α -sarcoglycan shRNA (h) Lentiviral Particles: sc-43416-V and α -sarcoglycan shRNA (m) Lentiviral Particles: sc-43417-V.


Molecular Weight of α -sarcoglycan: 50 kDa.

Positive Controls: α -sarcoglycan (h): 293T Lysate: sc-114140.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



α -sarcoglycan (C-15): sc-14174. Western blot analysis of α -sarcoglycan expression in non-transfected: sc-117750 (A) and human α -sarcoglycan transfected: sc-114140 (B) whole cell lysates.



α -sarcoglycan (C-15): sc-14174. Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing membrane and cytoplasmic staining of myocytes.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.