

# CYP19 (C-16): sc-14245

## BACKGROUND

P450 enzymes constitute a family of monooxygenase enzymes that are involved in the metabolism of a wide array of endogenous and xenobiotic compounds. P450 enzymes can be classified, based on their sequence similarities, into distinct subfamilies, which include CYP1A and CYP2A. Other P450 family members include CYP19, also designated aromatase (P450arom), which catalyzes the conversion of C19 steroids to estrogens in various tissues, including placenta, gonads, adipose tissue, skin and brain. CYP19 expression is controlled by hormonally regulated promoters in different tissues and increased aromatase activity is associated with familial gynecomastia. Also, a polymorphic allele of CYP19 (repeat (TTTA)<sub>12</sub>) is present in a majority of breast cancer patients. P450 cholesterol 7 $\alpha$ -hydroxylase, CYP7A1, is the rate limiting enzyme of bile acid synthesis in the liver, and its expression is mediated by the bile acid receptor FXR. CYP27A1 catalyzes vitamin D3 25-hydroxylation and is localized to the mitochondria in kidney and liver.

## CHROMOSOMAL LOCATION

Genetic locus: CYP19A1 (human) mapping to 15q21.2; Cyp19a1 (mouse) mapping to 9 A5.3.

## SOURCE

CYP19 (C-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of CYP19 of human origin.

## PRODUCT

Each vial contains 100  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-14245 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

CYP19 (C-16) is recommended for detection of CYP19 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CYP19 (C-16) is also recommended for detection of CYP19 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for CYP19 siRNA (h): sc-41498, CYP19 siRNA (m): sc-41499, CYP19 shRNA Plasmid (h): sc-41498-SH, CYP19 shRNA Plasmid (m): sc-41499-SH, CYP19 shRNA (h) Lentiviral Particles: sc-41498-V and CYP19 shRNA (m) Lentiviral Particles: sc-41499-V.

Molecular Weight (predicted) of CYP19: 58 kDa.

Molecular Weight (observed) of CYP19: 50 kDa.


## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## DATA



CYP19 (C-16): sc-14245. Western blot analysis of CYP19 expression in rat ovary tissue extract.

## SELECT PRODUCT CITATIONS

- Jesmin, S., et al. 2004. Aromatase is abundantly expressed by neonatal rat penis but downregulated in adulthood. *J. Mol. Endocrinol.* 33: 343-359.
- Wagner, S., et al. 2008. Cytochrome b5 expression in gonadectomy-induced adrenocortical neoplasms of the domestic ferret (*Mustela putorius furo*). *Vet. Pathol.* 45: 439-442.
- Luczak, M.W. and Jagodzinski, P.P. 2008. Trichostatin A down-regulates CYP19 transcript and protein levels in MCF-7 breast cancer cells. *Biomed. Pharmacother.* 63: 262-266.
- Gross, M., et al. 2009. Expression of androgen and estrogen related proteins in normal weight and obese prostate cancer patients. *Prostate* 69: 520-527.
- Han, J., et al. 2009. Increased expression of the testicular estrogen receptor  $\alpha$  in adult mice exposed to low doses of methiocarb. *J. Appl. Toxicol.* 29: 446-451.
- Zhao, Y., et al. 2010. Perfluorooctanoic acid effects on steroid hormone and growth factor levels mediate stimulation of peripubertal mammary gland development in C57BL/6 mice. *Toxicol. Sci.* 115: 214-224.
- Sarachana, T., et al. 2011. Sex hormones in autism: androgens and estrogens differentially and reciprocally regulate RORA, a novel candidate gene for autism. *PLoS ONE* 6: e17116.
- Pöllänen, E., et al. 2011. Differential influence of peripheral and systemic sex steroids on skeletal muscle quality in pre- and postmenopausal women. *Aging Cell* 10: 650-660.
- Wang, W., et al. 2012. Cortisol induces aromatase expression in human placental syncytiotrophoblasts through the cAMP/Sp1 pathway. *Endocrinology* 153: 2012-2022.

**MONOS**  
Satisfaction  
Guaranteed

Try **CYP19 (E-9): sc-374176**, our highly recommended monoclonal alternative to CYP19 (C-16). Also, for AC, HRP, FITC, PE, Alexa Fluor<sup>®</sup> 488 and Alexa Fluor<sup>®</sup> 647 conjugates, see **CYP19 (E-9): sc-374176**.