

OPN1SW (H-17): sc-14365

BACKGROUND

G protein-coupled receptors (GPCRs), which are characterized by containing seven transmembrane α helices, elicit G protein-mediated signaling cascades in response to a variety of stimuli. The opsin subfamily, which represents approximately 90 percent of all GPCRs, is comprised of photoreceptors that are activated by light. It includes the red, green and blue-sensitive opsins and rhodopsin. The opsin subfamily consists of an apoprotein covalently linked to 11-*cis*-retinal, which undergoes isomerization upon the absorption of photons. This isomerization leads to a conformational change of the protein, which results in the activation of hundreds of G proteins. Color is perceived in humans by three pigments, which localize to retinal cone photoreceptor cells. They are the blue-, green- and red-sensitive opsins, which are encoded by OPN1SW, OPN1MW and OPN1LW, respectively. Mutations in the OPN1MW and OPN1LW encoded opsins lead to the X-linked disorders protanopia and deuteranopia, respectively. Mutations in the OPN1SW encoded opsin leads to tritanopia, an autosomal dominant disorder, which is characterized by decreased sensitivity to blue light.

REFERENCES

1. Fung, B.K., et al. 1980. Flow of information in the light-triggered cyclic nucleotide cascade of vision. *Proc. Natl. Acad. Sci. USA* 78: 152-156.
2. Hargrave, P.A., et al. 1983. The structure of bovine rhodopsin. *Biophys. Struct. Mech.* 9: 235-244.
3. Drummond-Borg, M., et al. 1988. Molecular basis of abnormal red-green color vision: a family with three types of color vision defects. *Am. J. Hum. Genet.* 43: 675-683.
4. Oprian, D.D., et al. 1991. Design, chemical synthesis, and expression of genes for the three human color vision pigments. *Biochemistry* 30: 11367-11372.

CHROMOSOMAL LOCATION

Genetic locus: OPN1SW (human) mapping to 7q32.1; Opn1sw (mouse) mapping to 6 A3.3.

SOURCE

OPN1SW (H-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of the opsin protein encoded by OPN1SW of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-14365 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

OPN1SW (H-17) is recommended for detection of the opsin protein encoded by OPN1SW of mouse, rat and, to a lesser extent, human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

OPN1SW (H-17) is also recommended for detection of the opsin protein encoded by OPN1SW in additional species, including canine.

Suitable for use as control antibody for OPN1SW siRNA (h): sc-40142, OPN1SW siRNA (m): sc-40143, OPN1SW shRNA Plasmid (h): sc-40142-SH, OPN1SW shRNA Plasmid (m): sc-40143-SH, OPN1SW shRNA (h) Lentiviral Particles: sc-40142-V and OPN1SW shRNA (m) Lentiviral Particles: sc-40143-V.


Molecular Weight of OPN1SW: 40 kDa.

Positive Controls: mouse eye extract: sc-364241.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



OPN1SW (H-17): sc-14365. Western blot analysis of OPN1SW expression in mouse eye tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.