

Synaptotagmin III (E-17): sc-14443

BACKGROUND

Synaptotagmins are a large gene family of synaptic vesicle type III integral membrane proteins that function as regulators of both exocytosis and endocytosis and are involved in neurotransmitter secretion from small secretory vesicles. Calcium binds to Synaptotagmin I which triggers neurotransmitter release at the synapse. Synaptotagmin II is phosphorylated by WNK1 in a process that regulates calcium-dependent interactions. Synaptotagmin III is involved in calcium-dependent exocytosis of secretory vesicles in endocrine cells and neurons. Synaptotagmin IV is expressed in neuronal tissues, and has the highest mRNA levels in the hippocampus. The proximity of the Synaptotagmin IV gene to markers of several psychiatric disorders suggest an involvement of Synaptotagmin IV in human disease. Synaptotagmin V is a dense-core vesicle-specific protein that regulates a specific type of calcium-regulated secretion. Synaptotagmin VI interacts with adaptor protein-2 in a calcium-independent manner. Synaptotagmin VII is widely expressed in non-neuronal tissues.

REFERENCES

- Hilbush, B.S. and Morgan, J.I. 1994. A third Synaptotagmin gene, Syt3, in the mouse. *Proc. Natl. Acad. Sci. USA* 91: 8195-8199.
- Li, C., et al. 1995. Ca²⁺-dependent and -independent activities of neural and non-neural Synaptotagmins. *Nature* 375: 594-599.
- Kishore, B.K., et al. 1998. Expression of Synaptotagmin VIII in rat kidney. *Am. J. Physiol.* 275: 131-142.
- Xi, D., et al. 1999. Analysis of Synaptotagmin I-IV messenger RNA expression and developmental regulation in the rat hypothalamus and pituitary. *Neuroscience* 88: 425-435.
- Ferguson, G.D., et al. 2000. The human Synaptotagmin IV gene defines an evolutionary break point between syntenic mouse and human chromosome regions but retains ligand inducibility and tissue specificity. *J. Biol. Chem.* 275: 36920-3696.

CHROMOSOMAL LOCATION

Genetic locus: Syt3 (mouse) mapping to 7 B4.

SOURCE

Synaptotagmin III (E-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Synaptotagmin III of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-14443 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Synaptotagmin III (E-17) is recommended for detection of Synaptotagmin III of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Synaptotagmin III (E-17) is also recommended for detection of Synaptotagmin III in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Synaptotagmin III siRNA (m): sc-41315, Synaptotagmin III shRNA Plasmid (m): sc-41315-SH and Synaptotagmin III shRNA (m) Lentiviral Particles: sc-41315-V.


Molecular Weight of Synaptotagmin III: 80 kDa.

Positive Controls: mouse cerebellum extract: sc-2403 or mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Synaptotagmin III (E-17): sc-14443. Western blot analysis of Synaptotagmin III expression in mouse cerebellum (A) and mouse brain (B) tissue extracts.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.