

PDK3 (A-16): sc-14490

BACKGROUND

Pyruvate dehydrogenase kinase family members (PDK1, 2, 3 and 4) are Serine kinases that catalyze phosphorylation of the E1 α subunit of the pyruvate dehydrogenase complex (PDC). PDC activity is controlled through phosphorylation and dephosphorylation of the E1 α subunit, which leads to inactivation and reactivation, respectively. PDK3 binding to a free lipoyl domain (L2) in dihydrolypoyl acetyltransferase (E2), which comprises the core of PDC, leads to a large increase in E1 α phosphorylation. Upregulation of PDK isoenzymes occurs during starvation conditions, where acetyl-CoA is alternatively generated through fatty acid oxidation. PDKs contain five conserved regions and are mechanistically similar to bacterial His-kinases in that both require histidine residues for activity. In mammals, transcripts for PDK3 are most abundant in testis and are moderately expressed in heart and skeletal muscle.

REFERENCES

- Gudi, R., et al. 1995. Diversity of the pyruvate dehydrogenase kinase gene family in humans. *J. Biol. Chem.* 270: 28989-28994.
- Bowker-Kinley, M.M., et al. 1998. Evidence for existence of tissue-specific regulation of the mammalian pyruvate dehydrogenase complex. *Biochem. J.* 329: 191-196.
- Sugden, M.C., et al. 2000. Selective modification of the pyruvate dehydrogenase kinase isoform profile in skeletal muscle in hyperthyroidism: implications for the regulatory impact of glucose on fatty acid oxidation. *J. Endocrinol.* 167: 339-345.
- Mooney, B.P., et al. 2000. Histidine modifying agents abolish pyruvate dehydrogenase kinase activity. *Biochem. Biophys. Res. Commun.* 267: 500-503.
- Baker, J.C., et al. 2000. Marked differences between two isoforms of human pyruvate dehydrogenase kinase. *J. Biol. Chem.* 275: 15773-15781.
- Wu, P., et al. 2000. Starvation increases the amount of pyruvate dehydrogenase kinase in several mammalian tissues. *Arch. Biochem. Biophys.* 381: 1-7.

CHROMOSOMAL LOCATION

Genetic locus: PDK3 (human) mapping to Xp22.11.

SOURCE

PDK3 (A-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PDK3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-14490 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4 $^{\circ}$ C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

PDK3 (A-16) is recommended for detection of PDK3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PDK3 (A-16) is also recommended for detection of PDK3 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for PDK3 siRNA (h): sc-39029, PDK3 shRNA Plasmid (h): sc-39029-SH and PDK3 shRNA (h) Lentiviral Particles: sc-39029-V.


Molecular Weight of PDK3: 47 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206 or PDK3 (h): 293T Lysate: sc-158838.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



PDK3 (A-16): sc-14490. Western blot analysis of PDK3 expression in non-transfected: sc-110760 (A) and human PDK3 transfected: sc-158838 (B) 293 whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **PDK3 (A-4): sc-365378** or **PDK3 (RR-2): sc-100535**, our highly recommended monoclonal alternatives to PDK3 (A-16).