

MCH-1R (W-17): sc-14511

BACKGROUND

Melanin-concentrating hormone (MCH) is a 19 amino acid cyclic neuropeptide, which is mainly expressed in the hypothalamus. MCH modulates feeding behavior, aggression, anxiety, arousal and reproductive function in mammals by controlling the release of luteinizing hormone (LH). The melanin-concentrating hormone receptor (MCHR, also designated SLC-1) is a glycosylated G-protein coupled receptor. MCHR mediates the effects of MCH through Gai and/or Gaq signaling and is expressed in several regions of the brain, including the cerebral cortex, amygdala, thalamus and hypothalamus (1,2,6). MCH and MCHR have also been implicated in stimulating leptin expression and secretion in adipocytes, which suggests that the melanin-concentrating hormone and its receptor may be potential targets for modulating obesity.

REFERENCES

1. Drozd, R., Hintermann, E., Tanner, H., Zumsteg, U. and Eberle, A.N. 1999. (D-(p-benzoylphenylalanine) 13, tyrosine19)-melanin-concentrating hormone, a potent analogue for MCH receptor crosslinking. *J. Pept. Sci.* 5: 234-242.
2. Saito, Y., Nothacker, H.P., Wang, Z., Lin, S.H., Leslie, F., and Civelli, O. 1999. Molecular characterization of the melanin-concentrating-hormone receptor. *Nature* 400: 265-269.
3. Murray, J.F., Baker, B.I., Levy, A. and Wilson, C.A. 2000. The influence of gonadal steroids on pre-pro melanin-concentrating hormone mRNA in female rats. *J. Neuroendocrinol.* 12: 53-59.
4. Murray, J.F., Adan, R.A., Walker, R., Baker, B.I., Thody, A.J., Nijenhuis, W.A., Yukitake, J. and Wilson, C.A. 2000. Melanin-concentrating hormone, melanocortin receptors and regulation of luteinizing hormone release. *J. Neuroendocrinol.* 12: 217-223.
5. Hervieu, G.J., Cluderay, J.E., Harrison, D., Meakin, J., Maycox, P., Nasir, S. and Leslie, R.A. 2000. The distribution of the mRNA and protein products of the melanin-concentrating hormone (MCH) receptor gene, slc-1, in the central nervous system of the rat. *Eur. J. Neurosci.* 12: 1194-1216.
6. Hawes, B.E., Kil, E., Green, B., O'Neill, K., Fried, S. and Graziano, M.P. 2000. The melanin-concentrating hormone receptor couples to multiple G proteins to activate diverse intracellular signaling pathways. *Endocrinology* 141: 4524-4532.

CHROMOSOMAL LOCATION

Genetic locus: MCHR1 (human) mapping to 22q13.2.

SOURCE

MCH-1R (W-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MCH-1R of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-14511 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MCH-1R (W-17) is recommended for detection of MCH-1R of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MCH-1R siRNA (h): sc-42017, MCH-1R shRNA Plasmid (h): sc-42017-SH and MCH-1R shRNA (h) Lentiviral Particles: sc-42017-V.

Molecular Weight of MCH-1R: 53 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **MCH-1R (52-W7): sc-100327**, our highly recommended monoclonal alternative to MCH-1R (W-17).