

c-Kit (M-14): sc-1494

BACKGROUND

The c-Kit proto-oncogene is a member of the receptor tyrosine kinase family and, more specifically, is closely related to the platelet derived growth factor receptor (PDGFR). c-Kit, the normal cellular homolog of the HZ4-feline sarcoma virus transforming gene (v-Kit), encodes a transmembrane receptor. c-Kit regulates a variety of biological responses including chemotaxis, cell proliferation, apoptosis and adhesion. c-Kit is also identical with the product of the W locus in mice and, as such, is integral to the development of mast cells and hematopoiesis. The ligand for the c-Kit receptor (KL) has been identified and is encoded at the murine steel (Sl) locus. Kit is the human homolog of the proto-oncogene c-Kit. Mutations in Kit are integral for tumor growth and progression in various cancers.

CHROMOSOMAL LOCATION

Genetic locus: KIT (human) mapping to 4q12; Kit (mouse) mapping to 5 C3.3.

SOURCE

c-Kit (M-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of c-Kit of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-1494 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

c-Kit (M-14) is recommended for detection of c-Kit p145 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

c-Kit (M-14) is also recommended for detection of c-Kit p145 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for c-Kit siRNA (h): sc-29225, c-Kit siRNA (m): sc-29852, c-Kit shRNA Plasmid (h): sc-29225-SH, c-Kit shRNA Plasmid (m): sc-29852-SH, c-Kit shRNA (h) Lentiviral Particles: sc-29225-V and c-Kit shRNA (m) Lentiviral Particles: sc-29852-V.

Molecular Weight of c-Kit precursor: 120 kDa.

Molecular Weight of mature c-Kit: 145 kDa.

Positive Controls: CCRF-HSB-2 cell lysate: sc-2265, HEL 92.1.7 cell lysate: sc-2270 or TF-1 cell lysate: sc-2412.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



c-Kit (M-14): sc-1494. Western blot analysis of c-Kit expression in CCRF-HSB-2 (A), HEL 92.1.7 (B) and TF-1 (C) whole cell lysates.



c-Kit (M-14): sc-1494. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

SELECT PRODUCT CITATIONS

- Schrans-Stassen, B., et al. 1999. Differential expression of c-Kit in mouse undifferentiated and differentiating type A spermatogonia. *Endocrinology* 140: 5894-5900.
- Wang, X.Y., et al. 2007. Unique distribution of interstitial cells of Cajal in the feline pylorus. *Cell Tissue Res.* 329: 13-24.
- Cervelló, I., et al. 2007. Identification, characterization and co-localization of label-retaining cell population in mouse endometrium with typical undifferentiated markers. *Hum Reprod.* 22: 45-51.
- Ishiguro, K., et al. 2009. Inhibition of all-trans retinoic acid-induced granulocytic differentiation of WEHI-3B D+ cells by forced expression of SCL (TAL1) and GATA-1. *Leuk. Res.* 33: 1249-1254.
- Drube, S., et al. 2010. The receptor tyrosine kinase c-Kit controls IL-33 receptor signaling in mast cells. *Blood* 115: 3899-3906.
- Wiebe, M.S., et al. 2010. Mice deficient in the serine/threonine protein kinase VRK1 are infertile due to a progressive loss of spermatogonia. *Biol. Reprod.* 82: 182-193.
- Deneubourg, L., et al. 2011. Abnormal elevated PTEN expression in the mouse antrum of a model of GIST Kit(K641E/K641E). *Cell. Signal.* 23: 1857-1868.
- Kurth, T.B., et al. 2011. Functional mesenchymal stem cell niches in the adult knee joint synovium *in vivo*. *Arthritis Rheum.* 63: 1289-1300.
- Gromova, P., et al. 2012. ENDOGLIN/CD105 is expressed in KIT positive cells in the gut and in gastrointestinal stromal tumours. *J. Cell. Mol. Med.* 16: 306-317.



Try **c-Kit (E-3): sc-365504** or **c-Kit (Ab 81): sc-13508**, our highly recommended monoclonal alternatives to c-Kit (M-14). Also, for AC, HRP, FITC, PE, Alexa Fluor[®] 488 and Alexa Fluor[®] 647 conjugates, see **c-Kit (E-3): sc-365504**.