

creatine kinase-M (C-14): sc-15164

BACKGROUND

Creatine kinases (CKs) are a large family of isoenzymes that regulate levels of ATP in subcellular compartments, where they provide ATP at sites of fluctuating energy demand by the transfer of phosphates between creatine and adenine nucleotides. Creatine kinases provide the energy of phosphate hydrolysis necessary to drive the normal function of many cellular systems including muscle, electrocytes, retina photoreceptor cells, brain cells, kidney, salt glands, myometrium, placenta, pancreas, thymus, thyroid, intestinal epithelial cells, endothelial cells, cartilage and bone cells, macrophages, blood platelets, and tumor and cancer cells. Human cytoplasmic creatine kinase-B, also designated CK-B and BCK, is a 381 amino acid, brain tissue-specific isoform of creatine kinase. Human cytoplasmic creatine kinase-M (CK-M, MCK) is a muscle tissue-specific isoform of creatine kinase. Human cytoplasmic creatine kinase-Mi (Mi-CK, MtCK) is a 416 amino acid mitochondrial-specific isoform of creatine kinase. Cytosolic creatine kinases are important in the energetic regulation of Ca^{2+} -pumps and in the maintenance of Ca^{2+} -homeostasis.

REFERENCES

1. Mariman, E.C., et al. 1987. Structure and expression of the human creatine kinase B gene. *Genomics* 1: 126-137.
2. Nigro, J.M., et al. 1987. cDNA cloning and mapping of the human creatine kinase M gene to 19q13. *Am. J. Hum. Genet.* 40: 115-125.
3. Haas, R.C., et al. 1989. Isolation and characterization of the gene and cDNA encoding human mitochondrial creatine kinase. *J. Biol. Chem.* 264: 2890-2897.
4. Mariman, E.C., et al. 1989. Complete nucleotide sequence of the human creatine kinase B gene. *Nucleic Acids Res.* 17: 6385.

CHROMOSOMAL LOCATION

Genetic locus: CKM (human) mapping to 19q13.32; Ckm (mouse) mapping to 7 A3.

SOURCE

creatine kinase-M (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of creatine kinase-M of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-15164 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

creatine kinase-M (C-14) is recommended for detection of creatine kinase-M chain of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


creatine kinase-M (C-14) is also recommended for detection of creatine kinase-M chain in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for creatine kinase-M siRNA (h): sc-35109, creatine kinase-M siRNA (m): sc-35110, creatine kinase-M shRNA Plasmid (h): sc-35109-SH, creatine kinase-M shRNA Plasmid (m): sc-35110-SH, creatine kinase-M shRNA (h) Lentiviral Particles: sc-35109-V, and creatine kinase-M shRNA (m) Lentiviral Particles: sc-35110-V.


Molecular Weight of creatine kinase-M: 43 kDa.

Positive Controls: human skeletal muscle extract: sc-363776, human heart extract: sc-363763 or HeLa whole cell lysate: sc-2200.

DATA



creatine kinase-M (C-14): sc-15164. Western blot analysis of creatine kinase-M expression in human skeletal muscle (A) and human heart (B) tissue extracts.



creatine kinase-M (C-14): sc-15164. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic staining of myocytes (B).

SELECT PRODUCT CITATIONS

1. Lees, S.J., et al. 2006. Age-associated decrease in muscle precursor cell differentiation. *Am. J. Physiol., Cell Physiol.* 290: C609-C615.
2. Graciotti, L., et al. 2011. Dystrophin is required for the normal function of the cardio-protective K(ATP) channel in cardiomyocytes. *PLoS ONE* 6: e27034.

MONOS
Satisfaction
Guaranteed

Try **creatine kinase-M (G-9): sc-365046** or **creatine kinase-M (H-9): sc-365141**, our highly recommended monoclonal alternatives to creatine kinase-M (C-14).