

UPIIIa (M-17): sc-15186

BACKGROUND

The asymmetric unit membrane (AUM) forms numerous plaques, which cover the apical surface of the urothelium. These plaques are thought to strengthen the urothelium and reduce the risk of rupturing during ladder distention. They are composed of four major integral membrane proteins called uroplakins (UP). The uroplakin family comprises UPIa, UPIb, UPII, and UPIII. Family members are conserved among several species, including human, mouse, rat, rabbit, canine, porcine and ovine. UPIa and UPIb form tightly packed structures with UPII and UPIII, respectively. This pairing is required for normal urothelial plaque formation and is regulated by proteolytic processing of the uroplakin proteins. Uroplakins are expressed in normal urothelium and are used as specific markers of urothelial differentiation. They are also expressed in a majority of transitional cell carcinomas of the bladder (TCCs), which make the uroplakins a useful marker for detecting bladder cancer metastasis and for staging and monitoring chemotherapeutic response.

REFERENCES

- Lin, J.H., et al. 1994. Precursor sequence, processing, and urothelium-specific expression of a major 15 kDa protein subunit of asymmetric unit membrane. *J. Biol. Chem.* 269: 1775-1784.
- Wu, X.R., et al. 1994. Mammalian uroplakins. A group of highly conserved urothelial differentiation-related membrane proteins. *J. Biol. Chem.* 269: 13716-13724.
- Wu, X.R., et al. 1995. Selective interactions of UPIa and UPIb, two members of the transmembrane 4 superfamily, with distinct single transmembrane-domained proteins in differentiated urothelial cells. *J. Biol. Chem.* 270: 29752-29759.

CHROMOSOMAL LOCATION

Genetic locus: UPK3A (human) mapping to 22q13.31; Upk3a (mouse) mapping to 15 E2.

SOURCE

UPIIIa (M-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of UPIIIa of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-15186 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

UPIIIa (M-17) is recommended for detection of UPIIIa of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

UPIIIa (M-17) is also recommended for detection of UPIIIa in additional species, including equine, canine and porcine.

Suitable for use as control antibody for UPIIIa siRNA (h): sc-41096, UPIIIa siRNA (m): sc-41097, UPIIIa shRNA Plasmid (h): sc-41096-SH, UPIIIa shRNA Plasmid (m): sc-41097-SH, UPIIIa shRNA (h) Lentiviral Particles: sc-41096-V and UPIIIa shRNA (m) Lentiviral Particles: sc-41097-V.


Molecular Weight of UPIIIa: 47 kDa.

Positive Controls: mouse bladder extract: sc-364919.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



UPIIIa (M-17): sc-15186. Western blot analysis of UPIIIa expression in mouse bladder tissue extract.



UPIIIa (M-17): sc-15186. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human urinary bladder tissue showing cytoplasmic and membrane staining of urothelial cells (B).

SELECT PRODUCT CITATIONS

- Jaal, J., et al. 2006. Radiation-induced damage to mouse urothelial barrier. *Radiother. Oncol.* 80: 250-256.