

P2Y6 (N-16): sc-15215

BACKGROUND

Nucleotides are emerging as important extracellular signaling molecules that mediate several effects, such as proliferation, differentiation, chemotaxis and cytokine release. The P2 receptor family is activated by the binding of nucleotides and is divided into two subfamilies, P2X and P2Y. The P2X receptor family is comprised of ligand-gated ion channels that allow for the increased permeability of calcium into the cell in response to extracellular ATP. The P2Y receptor family consists of eleven members of G protein-coupled receptors, P2Y1, P2Y2, P2Y3, P2Y4, P2Y5, P2Y6, P2Y7, P2Y8, P2Y9, P2Y10 and P2Y11, which mediate the effects of extracellular nucleotides, primarily through the activation of phospholipase C. To some extent, the P2Y receptors can also activate potassium channels or, alternatively, inhibit adenylate cyclase and N-type calcium channels in response to extracellular nucleotides. The P2Y receptors are differentially expressed in several tissue in leukocytes, which suggests a role for the P2Y receptor family in the activation of leukocytes and platelets in response to inflammation or vascular damage.

CHROMOSOMAL LOCATION

Genetic locus: P2RY6 (human) mapping to 11q13.4; P2ry6 (mouse) mapping to 7 E3.

SOURCE

P2Y6 (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of P2Y6 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-15215 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

P2Y6 (N-16) is recommended for detection of P2Y6 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

P2Y6 (N-16) is also recommended for detection of P2Y6 in additional species, including canine.


Suitable for use as control antibody for P2Y6 siRNA (h): sc-42584, P2Y6 siRNA (m): sc-42585, P2Y6 shRNA Plasmid (h): sc-42584-SH, P2Y6 shRNA Plasmid (m): sc-42585-SH, P2Y6 shRNA (h) Lentiviral Particles: sc-42584-V and P2Y6 shRNA (m) Lentiviral Particles: sc-42585-V.

Molecular Weight of P2Y6: 36 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



P2Y6 (N-16): sc-15215. Western blot analysis of P2Y6 expression in mouse PBL whole cell lysate.

SELECT PRODUCT CITATIONS

1. Yu, Q., et al. 2011. Expression of P2Y receptors in the rat anterior pituitary. *Purinergic Signal.* 7: 207-219.
2. Chen, D., et al. 2011. Expression of P2Y6 receptors in the developing mouse skeletal muscle and after injury and repair. *J. Anat.* 218: 643-651.
3. Yu, W., et al. 2013. Extracellular UDP enhances P2X-mediated bladder smooth muscle contractility via P2Y6 activation of the phospholipase C/inositol trisphosphate pathway. *FASEB J.* 27: 1895-1903.
4. Matta, C., et al. 2014. Purinergic signalling is required for calcium oscillations in migratory chondrogenic progenitor cells. *Pflugers Arch.* 467: 429-442.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.