

# Integrin $\beta 7$ (H-120): sc-15330

## BACKGROUND

Integrins are heterodimers composed of noncovalently associated transmembrane  $\alpha$  and  $\beta$  subunits. The  $16\alpha$  and  $8\beta$  subunits heterodimerize to produce more than 20 different receptors. Most Integrin receptors bind ligands that are components of the extracellular matrix, including fibronectin, collagen and vitronectin. Certain Integrins can also bind to soluble ligands such as fibrinogen, or to counterreceptors on adjacent cells, such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster Integrins by binding to adjacent Integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of Integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organization, Integrins function as signaling receptors. Signals transduced by Integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis.

## REFERENCES

- Hynes, R.O. 1992. Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* 69: 11-25.
- Miyamoto, S., Akiyama, S.K., Yamada, K.M. 1995. Synergistic roles for receptor occupancy and aggregation in Integrin transmembrane function. *Science* 267: 883-885.
- Clark, E.A. and Brugge, J.S. 1995. Integrins and signal transduction pathways: the road taken. *Science* 268: 233-239.
- Sheppard, D. 1996. Epithelial integrins. *BioEssays* 18: 655-660.
- Juliano, R. 1996. Cooperation between soluble factors and integrin-mediated cell anchorage in the control of cell growth and differentiation. *BioEssays* 18: 911-917.

## CHROMOSOMAL LOCATION

Genetic locus: ITGB7 (human) mapping to 12q13.13; Itgb7 (mouse) mapping to 15 F3.

## SOURCE

Integrin  $\beta 7$  (H-120) is a rabbit polyclonal antibody corresponding to amino acids 641-760 mapping near the C-terminus of Integrin  $\beta 7$  of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## APPLICATIONS

Integrin  $\beta 7$  (H-120) is recommended for detection of Integrin  $\beta 7$  of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Integrin  $\beta 7$  (H-120) is also recommended for detection of Integrin  $\beta 7$  in additional species, including equine, canine and porcine.

Suitable for use as control antibody for Integrin  $\beta 7$  siRNA (h): sc-35682, Integrin  $\beta 7$  siRNA (m): sc-37260, Integrin  $\beta 7$  shRNA Plasmid (h): sc-35682-SH, Integrin  $\beta 7$  shRNA Plasmid (m): sc-37260-SH, Integrin  $\beta 7$  shRNA (h) Lentiviral Particles: sc-35682-V and Integrin  $\beta 7$  shRNA (m) Lentiviral Particles: sc-37260-V.

Molecular Weight of Integrin  $\beta 7$ : 140 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, ECV304 cell lysate: sc-2269 or Ramos cell lysate: sc-2216.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/ 2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **Integrin  $\beta 7$  (D-5): sc-515397** or **Integrin  $\beta 7$  (C-2): sc-166031**, our highly recommended monoclonal alternatives to Integrin  $\beta 7$  (H-120).