

Atm (H-280): sc-15392

BACKGROUND

The phosphatidylinositol kinase (PIK) family members fall into two distinct subgroups. The first subgroup contains proteins such as the PI 3- and PI 4-kinases and the second group comprises the PIK-related kinases. The PIK-related kinases include Atm, DNA-PKCS and FRAP. These proteins have in common a region of homology at their carboxy-termini that is not present in the PI 3- and PI 4-kinases. The Atm gene is mutated in the autosomal recessive disorder ataxia telangiectasia (AT) that is characterized by cerebellar degeneration (ataxia) and the appearance of dilated blood vessels (telangiectases) in the conjunctivae of the eyes. AT cells are hypersensitive to ionizing radiation, impaired in mediating the inhibition of DNA synthesis and display delays in p53 induction.

REFERENCES

- Hartley, K.O., et al. 1995. DNA-dependent protein kinase catalytic subunit: a relative of phosphatidylinositol 3-kinase and the ataxia telangiectasia gene product. *Cell* 82: 849-856.
- Nowak, R. 1995. Discovery of AT gene sparks biomedical research bonanza. *Science* 268: 1700-1701.
- Savitsky, K., et al. 1995. A single ataxia telangiectasia gene with a product similar to PI-3 kinase. *Science* 268: 1749-1753.
- Keith, C.T., et al. 1995. PIK-related kinases: DNA repair, recombination, and cell cycle checkpoints. *Science* 270: 50-51.
- Hickson I., et al. 2004. Identification and characterization of a novel and specific inhibitor of the ataxia-telangiectasia mutated kinase Atm. *Cancer Res.* 64: 9152-9159.
- Chen L., et al. 2005. Ataxia-telangiectasia-mutated dependent phosphorylation of Artemis in response to DNA damage. *Cancer Sci.* 96: 134-141.
- Liu A., et al. 2005. Alterations of DNA damage-response genes Atm and ATR in pyothorax-associated lymphoma. *Lab. Invest.* 85: 436-446.
- Lavin M.F., et al. 2005. Atm signaling and genomic stability in response to DNA damage. *Mutat Res.* 569: 123-132.

CHROMOSOMAL LOCATION

Genetic locus: ATM (human) mapping to 11q22.3; Atm (mouse) mapping to 9 A5.3.

SOURCE

Atm (H-280) is a rabbit polyclonal antibody raised against amino acids 2581-2860 of Atm (ataxia telangiectasia protein) of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Atm (H-280) is recommended for detection of Atm of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Atm (H-280) is also recommended for detection of Atm in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Atm siRNA (h): sc-29761, Atm siRNA (m): sc-29762, Atm shRNA Plasmid (h): sc-29761-SH, Atm shRNA Plasmid (m): sc-29762-SH, Atm shRNA (h) Lentiviral Particles: sc-29761-V and Atm shRNA (m) Lentiviral Particles: sc-29762-V.

Molecular Weight of Atm: 370 kDa.

Positive Controls: KNRK nuclear extract: sc-2141, HeLa whole cell lysate: sc-2200 or RAW 264.7 whole cell lysate: sc-2211.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfation
Guaranteed

Try **Atm (G-12): sc-377293** or **Atm (1B10): sc-135663**, our highly recommended monoclonal alternatives to Atm (H-280). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **Atm (G-12): sc-377293**.