

Synaptotagmin V (N-17): sc-15492

BACKGROUND

Synaptotagmins are a large gene family of synaptic vesicle type III integral membrane proteins that function as regulators of both exocytosis and endocytosis and are involved in neurotransmitter secretion from small secretory vesicles. Calcium binds to Synaptotagmin I which triggers neurotransmitter release at the synapse. Synaptotagmin II is phosphorylated by WNK1 in a process that regulates calcium-dependent interactions. Synaptotagmin III is involved in calcium-dependent exocytosis of secretory vesicles in endocrine cells and neurons. Synaptotagmin IV is expressed in neuronal tissues, and has the highest mRNA levels in the hippocampus. The proximity of the Synaptotagmin IV gene to markers of several psychiatric disorders suggest an involvement of Synaptotagmin IV in human disease. Synaptotagmin V is a dense-core vesicle-specific protein that regulates a specific type of calcium-regulated secretion. Synaptotagmin VI interacts with adaptor protein-2 in a calcium-independent manner. Synaptotagmin VII is widely expressed in non-neuronal tissues.

REFERENCES

- Hilbush, B.S. and Morgan, J.I. 1994. A third Synaptotagmin gene, Syt3, in the mouse. *Proc. Natl. Acad. Sci. USA* 91: 8195-8199.
- Li, C., Ullrich, B., Zhang, J.Z., Anderson, R.G., Brose, N. and Sudhof, T.C. 1995. Ca²⁺-dependent and -independent activities of neural and non-neuronal synaptotagmins. *Nature* 375: 594-599.
- Kishore, B.K., Wade, J.B., Schorr, K., Inoue, T., Mandon, B. and Knepper, M.A. 1998. Expression of Synaptotagmin VIII in rat kidney. *Am. J. Physiol.* 275: 131-142.
- Xi, D., Chin, H. and Gainer, H. 1999. Analysis of Synaptotagmin I-IV messenger RNA expression and developmental regulation in the rat hypothalamus and pituitary. *Neuroscience* 88: 425-435.
- Ferguson, G.D., Chen, X.N., Korenberg, J.R. and Herschman, H.R. 2000. The human Synaptotagmin IV gene defines an evolutionary break point between syntenic mouse and human chromosome regions but retains ligand inducibility and tissue specificity. *J. Biol. Chem.* 275: 36920-3696.
- LocusLink Report (LocusID: 6860). <http://www.ncbi.nlm.nih.gov/>

CHROMOSOMAL LOCATION

Genetic locus: SYT5 (human) mapping to 19q13.42; Syt5 (mouse) mapping to 7 A1.

SOURCE

Synaptotagmin V (N-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of Synaptotagmin V of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-15492 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Synaptotagmin V (N-17) is recommended for detection of Synaptotagmin V of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Synaptotagmin V (N-17) is also recommended for detection of Synaptotagmin V in additional species, including bovine and porcine.

Suitable for use as control antibody for Synaptotagmin V siRNA (h): sc-41318, Synaptotagmin V siRNA (m): sc-41319, Synaptotagmin V shRNA Plasmid (h): sc-41318-SH, Synaptotagmin V shRNA Plasmid (m): sc-41319-SH, Synaptotagmin V shRNA (h) Lentiviral Particles: sc-41318-V and Synaptotagmin V shRNA (m) Lentiviral Particles: sc-41319-V.

Molecular Weight of Synaptotagmin V: 48 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Synaptotagmin V (F-7): sc-398837** or **Synaptotagmin V (46): sc-136090**, our highly recommended monoclonal alternatives to Synaptotagmin V (N-17).