



# atonal (dN-19): sc-15699

## BACKGROUND

*Drosophila melanogaster* is a proven and effective model for studying developmental and cellular processes common to higher eukaryotes. Approximately 13,600 genes have been elucidated from more than 120 megabases of euchromatin, and they are organized among the chromosomes 2, 3, 4, X and Y, with the Y chromosome being predominately heterochromatic. *Drosophila* genes can be categorized based on the type of protein for which they encode and are represented by six major classifications, which include intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing, and chromatin associated) or other functional proteins. Among these numerous proteins, atonal (CG7508, CT23037) is an RNA polymerase II (Pol II) transcription factor that is expressed in chordotonal organ progenitor cells and regulates brain development during embryogenesis.

## REFERENCES

1. Jarman, A.P., Grau, Y., Jan, L.Y., and Jan, Y.N. 1993. Atonal is a proneural gene that directs chordotonal organ formation in the *Drosophila* peripheral nervous system. *Cell* 73: 1307-1321.
2. Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D., Amanatides, P. et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* 287: 2185-2195.
3. zur Lage PI, Powell LM, Prentice DR, McLaughlin P, Jarman AP. 2004. EGF receptor signaling triggers recruitment of *Drosophila* sense organ precursors by stimulating proneural gene autoregulation. *Dev. Cell* 7: 687-696
4. The Interactive Fly. <http://sdb.bio.purdue.edu/fly/aimain/1aahome.htm>.  
<http://sdb.bio.purdue.edu/fly/neural/atonal.htm>
5. LocusLink Report (LocusID: 40975) <http://www.ncbi.nlm.nih.gov/LocusLink/>

## SOURCE

atonal (dN-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of atonal of *Drosophila melanogaster* origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-15699 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## APPLICATIONS

atonal (dN-19) is recommended for detection of atonal of *Drosophila melanogaster* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.