



## eyes absent (dE-20): sc-15754

### BACKGROUND

*Drosophila melanogaster* is a proven and effective model for studying developmental and cellular processes common to higher eukaryotes. Approximately 13,600 genes have been elucidated from more than 120 megabases of euchromatin, and they are organized among the chromosomes 2, 3, 4, X and Y, with the Y chromosome being predominately heterochromatic. *Drosophila* genes can be categorized based on the type of protein for which they encode and are represented by six major classifications, which include intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing, and chromatin associated) or other functional proteins. Among these numerous proteins, eyes absent (Clift, Cli, CG9554, CT27024, EY2-1, *eya*) is an essential factor for proper development of the compound eye and for the determination of somatic gonadal precursor (SGP) cell fate during embryogenesis.

### REFERENCES

1. Bonini, N.M., Leiserson, W.M., and Benzer, S. 1993. The eyes absent gene: genetic control of cell survival and differentiation in the developing *Drosophila* eye. *Cell* 72: 379-395.
2. Boyle, M., Bonini, N., and DiNardo, S. 1997. Expression and function of clift in the development of somatic gonadal precursors within the *Drosophila* mesoderm. *Development* 124: 971-982.
3. Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D., Amanatides, P., et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* 287: 2185-2195.
4. Zimmerman, J.E., Bui, Q.T., Liu, H., and Bonini, N.M. 2000. Molecular genetic analysis of *Drosophila* eyes absent mutants reveals an eye enhancer element. *Genetics* 154: 237-246.
5. The Interactive Fly. <http://www.sdbonline.org/fly/airmain/1aahome.htm>.  
<http://www.sdbonline.org/fly/newgene/eyesab1.htm>
6. LocusLink Report (LocusID: 33916). <http://www.ncbi.nlm.nih.gov/LocusLink/>

### SOURCE

eyes absent (dE-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of eyes absent of *Drosophila melanogaster* origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-15754 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### APPLICATIONS

eyes absent (dE-20) is recommended for detection of eyes absent of *Drosophila melanogaster* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.