Polycomb (dN-19): sc-15814



The Power to Question

BACKGROUND

Drosophila melanogaster is a proven and effective model for studying developmental and cellular processes common to higher eukaryotes. Approximately 13,600 genes have been elucidated from more than 120 megabases of euchromatin, and they are organized among the chromosomes 2, 3, 4, X and Y, with the Y chromosome being predominately heterochromatic. Drosophila genes can be categorized based on the type of protein for which they encode and are represented by six major classifications, which include intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing, and chromatin associated) or other functional proteins. Among these proteins, Polycomb (Pc) is a transcription regulator that prevents the interaction of transcriptional components with cis-elements by associating with cis-polycomb response elements (PREs) and creating novel chromatin structures.

REFERENCES

- Breiling, A., Bonte, E., Ferrari, S., Becker, P.B. and Paro, R. 1999. The Drosophila Polycomb protein interacts with nucleosomal core particles in vitro via its repression domain. Mol. Cell. Biol. 19: 8451-8460.
- Dietzel, S., Niemann, H., Bruckner, B., Maurange, C. and Paro, R. 1999.
 The nuclear distribution of Polycomb during *Drosophila melanogaster* development shown with a GFP fusion protein. Chromosoma 108: 83-94.
- 3. Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D., Amanatides, P., et al. 2000. The genome sequence of *Drosophila melanogaster*. Science 287: 2185-2195.
- Poux, S., McCabe, D. and Pirrotta, V. 2001. Recruitment of components of Polycomb group chromatin complexes in *Drosophila*. Development 128: 75-85.
- 5. The Interactive Fly. http://www.sdbonline.org/fly/aimain/1aahome.htm. http://www.sdbonline.org/fly/polycomb/polycmb1.htm
- 6. LocusLink Report (LocusID:40358). http://www.ncbi.nlm.nih.gov/LocusLink/

SOURCE

Polycomb (dN-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Polycomb of *Drosophila melanogaster* origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-15814 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Polycomb (dN-19) is recommended for detection of Polycomb of *Drosophila melanogaster* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1–2 μ g per 100–500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight: 64 kDa

Positive Controls: Schneider's Drosophilia line 2 whole cell lysate.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/ 2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Polycomb (dN-19): sc-15814. Western blot analysis of Polycomb expression in Schneider's *Drosophila* line 2 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com