



kismet (dK-20): sc-15848

BACKGROUND

Drosophila melanogaster is a proven and effective model for studying developmental and cellular processes common to higher eukaryotes. Approximately 13,600 genes have been elucidated from more than 120 megabases of euchromatin, and they are organized among the chromosomes 2, 3, 4, X and Y, with the Y chromosome being predominately heterochromatic. *Drosophila* genes can be categorized based on the type of protein they encode and are represented by six major classifications, which include intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing, and chromatin associated) or other functional proteins. Among these proteins, kismet (Kis) is a putative chromatin remodeling factor that is expressed along anterior-posterior axes and mediates proper larval body segmentation. The kismet gene encodes several large nuclear proteins.

REFERENCES

1. Daubresse, G., Deuring, R., Moore, L., Papoulas, O., Zakrajsek, I., Waldrip, W.R., Scott, M.P., Kennison, J.A. and Tamkun, J.W. 1999. The *Drosophila* kismet gene is related to chromatin-remodeling factors and is required for both segmentation and segment identity. *Development* 126: 1175-1187.
2. Therrien, M., Morrison, D.K., Wong, A.M. and Rubin, G.M. 2000. A genetic screen for modifiers of a kinase suppressor of Ras-dependent rough eye phenotype in *Drosophila*. *Genetics* 156: 1231-1242.
3. Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D., Amanatides, P., et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* 287: 2185-2195.
4. The Interactive Fly. <http://www.sdbonline.org/fly/aimain/1aahome.htm>.
<http://www.sdbonline.org/fly/aimain/6biochem.htm>.
5. FlyBase. [http://flybase.bio.indiana.edu/\(FlyBaseID:FBgn0001309\)](http://flybase.bio.indiana.edu/(FlyBaseID:FBgn0001309)).
6. LocusLink Report. [http://www.ncbi.nlm.nih.gov/LocusLink/\(LocusID:33185\)](http://www.ncbi.nlm.nih.gov/LocusLink/(LocusID:33185)).

SOURCE

kismet (dK-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of kismet of *Drosophila melanogaster* origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-15848 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

kismet (dK-20) is recommended for detection of kismet of *Drosophila melanogaster* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.