



# $\beta$ -1,4-GalNAc-T2 (K-17): sc-160000

## BACKGROUND

$\beta$ -1,4-N-acetyl-galactosaminyl transferase 2 ( $\beta$ -1,4-GalNAc-T2) is a 566 amino acid protein belonging to the glycosyltransferase 2 family. Localized to the membrane of the Golgi apparatus,  $\beta$ -1,4-GalNAc-T2 participates in the synthesis of the Sd(a) antigen, a carbohydrate determinant expressed on erythrocytes, colonic mucosa and other tissues. During Sd(a) production,  $\beta$ -1,4-GalNAc-T2 transfers a  $\beta$ -1,4-linked GalNAc to the galactose residue of an  $\alpha$ -2,3-sialylated chain.  $\beta$ -1,4-GalNAc-T2 also catalyzes the last step in the biosynthesis of the CAD antigen.  $\beta$ -1,4-GalNAc-T2 is widely expressed, with the highest expression in colon and lesser expression in kidney, stomach, ileum and rectum. Mutations in the gene encoding  $\beta$ -1,4-GalNAc-T2 have been linked to Type I von Willebrand disease (VWD), the most common bleeding disorder in humans, characterized by reduced levels of plasma von Willebrand factor. Two named isoforms of  $\beta$ -1,4-GalNAc-T2 exist as a result of alternative splicing events.

## REFERENCES

- Smith, P.L. et al. 1994. Molecular cloning of a murine N-acetylgalactosamine transferase cDNA that determines expression of the T lymphocyte-specific CT oligosaccharide differentiation antigen. *J. Biol. Chem.* 269: 15162-15171.
- Dohi, T., et al. 1996. Detection of N-acetylgalactosaminyltransferase mRNA which determines expression of Sda blood group carbohydrate structure in human gastrointestinal mucosa and cancer. *Int. J. Cancer.* 67: 626-631.
- Mohlke, K.L., et al. 1996. A novel modifier gene for plasma von Willebrand factor level maps to distal mouse chromosome 11. *Proc. Natl. Acad. Sci. USA* 93: 15352-15357.
- Hakomori, S. 1999. Antigen structure and genetic basis of histo-blood groups A, B and O: their changes associated with human cancer. *Biochim. Biophys. Acta* 1473: 247-266.
- Mohlke, K.L., et al. 1999. Mvwf, a dominant modifier of murine von Willebrand factor, results from altered lineage-specific expression of a glycosyltransferase. *Cell* 96: 111-120.
- Montiel, M.D., et al. 2003. Molecular cloning, gene organization and expression of the human UDP-GalNAc:Neu5Ac $\alpha$ 2-3Gal $\beta$ -R  $\beta$ -1,4-N-acetylgalactosaminyltransferase responsible for the biosynthesis of the blood group Sda/CAD antigen: evidence for an unusual extended cytoplasmic domain. *Biochem. J.* 373: 369-379.
- Lo Presti, L., et al. 2003. Molecular cloning of the human  $\beta$ -1,4 N-acetylgalactosaminyltransferase responsible for the biosynthesis of the Sd(a) histo-blood group antigen: the sequence predicts a very long cytoplasmic domain. *J. Biochem.* 134: 675-682.
- Sato, T., et al. 2003. Molecular cloning and characterization of a novel human  $\beta$ -1,4-N-acetylgalactosaminyltransferase,  $\beta$  4GalNAc-T3, responsible for the synthesis of N,N'-diacetyllactosdiamine, GalNAc  $\beta$ -1-4GlcNAc. *J. Biol. Chem.* 278: 47534-47544.

## CHROMOSOMAL LOCATION

Genetic locus: B4galnt2 (mouse) mapping to 11 D.

## SOURCE

$\beta$ -1,4-GalNAc-T2 (K-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of  $\beta$ -1,4-GalNAc-T2 of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-160000 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

$\beta$ -1,4-GalNAc-T2 (K-17) is recommended for detection of  $\beta$ -1,4-GalNAc-T2 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with family members  $\beta$ -1,4-GalNAc-T,  $\beta$ -1,4-GalNAc-T3, or  $\beta$ -1,4-GalNAc-T4.

Suitable for use as control antibody for  $\beta$ -1,4-GalNAc-T2 siRNA (m): sc-108229,  $\beta$ -1,4-GalNAc-T2 shRNA Plasmid (m): sc-108229-SH and  $\beta$ -1,4-GalNAc-T2 shRNA (m) Lentiviral Particles: sc-108229-V.

Molecular Weight of  $\beta$ -1,4-GalNAc-T2: 63 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.