



CBARA1 (D-13): sc-160210

BACKGROUND

The EF-hand domain is a 12 amino acid loop motif that is commonly found in proteins that participate in calcium binding events within the cell. EF-hand domains generally exist in a pair that together form a stable four-helix bundle that enables the binding of calcium ions. CBARA1 (calcium binding atopy-related autoantigen 1), also known as CALC or EFHA3, is a 476 amino acid single-pass membrane protein that contains two EF-hand domains. Expressed at high levels in epidermal keratinocytes and dermal endothelial cells, CBARA1 functions to induce T cell-mediated autoreactivity, which is accompanied by the release of IFN- γ and can induce an allergic reaction that leads to the formation of IgE. IgE can bind to otherwise innocuous environmental particles and, upon binding, can induce cross-linking with an IgE receptor, an event that is associated with atopic dermatitis (AD). Multiple isoforms of CBARA1 exist due to alternative splicing events.

REFERENCES

1. Moncrief, N.D., Kretsinger, R.H. and Goodman, M. 1990. Evolution of EF-hand calcium-modulated proteins. I. Relationships based on amino acid sequences. *J. Mol. Evol.* 30: 522-562.
2. Nakayama, S., Moncrief, N.D. and Kretsinger, R.H. 1992. Evolution of EF-hand calcium-modulated proteins. II. Domains of several subfamilies have diverse evolutionary histories. *J. Mol. Evol.* 34: 416-448.
3. Maruyama, K. and Sugano, S. 1994. Oligo-capping: a simple method to replace the cap structure of eukaryotic mRNAs with oligoribonucleotides. *Gene* 138: 171-174.
4. Kawasaki, H. and Kretsinger, R.H. 1995. Calcium-binding proteins 1: EF-hands. *Protein Profile* 2: 297-490.
5. Natter, S., Seiberler, S., Hufnagl, P., Binder, B.R., Hirschl, A.M., Ring, J., Abeck, D., Schmidt, T., Valent, P. and Valenta, R. 1998. Isolation of cDNA clones coding for IgE autoantigens with serum IgE from atopic dermatitis patients. *FASEB J.* 12: 1559-1569.
6. Online Mendelian Inheritance in Man, OMIM™. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 605084. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Aichberger, K.J., Mittermann, I., Reininger, R., Seiberler, S., Swoboda, I., Spitzauer, S., Kopp, T., Stingl, G., Sperr, W.R., Valent, P., Repa, A., Bohle, B., Kraft, D. and Valenta, R. 2005. Hom s 4, an IgE-reactive autoantigen belonging to a new subfamily of calcium-binding proteins, can induce Th cell type 1-mediated autoreactivity. *J. Immunol.* 175: 1286-1294.

CHROMOSOMAL LOCATION

Genetic locus: CBARA1 (human) mapping to 10q22.1; Cbara1 (mouse) mapping to 10 B4.

SOURCE

CBARA1 (D-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CBARA1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-160210 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CBARA1 (D-13) is recommended for detection of CBARA1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CBARA1 siRNA (h): sc-90788, CBARA1 siRNA (m): sc-142033, CBARA1 shRNA Plasmid (h): sc-90788-SH, CBARA1 shRNA Plasmid (m): sc-142033-SH, CBARA1 shRNA (h) Lentiviral Particles: sc-90788-V and CBARA1 shRNA (m) Lentiviral Particles: sc-142033-V.

Molecular Weight of CBARA1: 54 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.