

Na⁺ CP type Iα (C-18): sc-16031

BACKGROUND

Voltage-gated sodium channels are selective ion channels that regulate the permeability of sodium ions in excitable cells. During the propagation of an action potential, sodium channels allow an influx of sodium ions, which rapidly depolarize the cell. The three glycoproteins that comprise the voltage-gated sodium channel proteins include a pore-forming α subunit, a non-covalently associated $\beta 1$ subunit and a disulfide-linked $\beta 2$ subunit. The two β subunits regulate the level of channel expression, modulate gating and function as cell adhesion molecules for cellular aggregation and cytoskeleton interaction. The α subunits of sodium channels type I and III are predominantly expressed in neuronal cell bodies and proximal processes, while type II α subunits are more abundant along axons. The $\beta 1$ subunit of sodium channel type I is expressed in brain, skeletal and cardiac muscle. In the brain, $\beta 1$ and $\beta 2$ are highly expressed in Purkinje cells, and $\beta 1$ is also expressed in the pyramidal cells of the deep cerebellar nuclei. Impaired voltage-gated sodium channels lead to a number of diseases including myotonia.

REFERENCES

- Rosenfeld, J., Sloan-Brown, K. and George, A.L., Jr. 1997. A novel muscle sodium channel mutation causes painful congenital myotonia. *Ann. Neurol.* 42: 811-814.
- Catterall, W.A. 1999. Molecular properties of brain sodium channels: an important target for anticonvulsant drugs. *Adv. Neurol.* 79: 441-456.
- Whitaker, W.R., Clare, J.J., Powell, A.J., Chen, Y.H., Faull, R.L. and Emson, P.C. 2000. Distribution of voltage-gated sodium channel α subunit and β subunit mRNAs in human hippocampal formation, cortex, and cerebellum. *J. Comp. Neurol.* 422: 123-139.

CHROMOSOMAL LOCATION

Genetic locus: SCN1A (human) mapping to 2q24.3; Scn1a (mouse) mapping to 2 C1.3.

SOURCE

Na⁺ CP type Iα (C-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Na⁺ CP type Iα of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16031 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Na⁺ CP type Iα (C-18) is recommended for detection of sodium channel type Iα of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Na⁺ CP type Iα (C-18) is also recommended for detection of sodium channel type Iα in additional species, including bovine.

Suitable for use as control antibody for Na⁺ CP type Iα siRNA (h): sc-42642, Na⁺ CP type Iα siRNA (m): sc-42643, Na⁺ CP type Iα shRNA Plasmid (h): sc-42642-SH, Na⁺ CP type Iα shRNA Plasmid (m): sc-42643-SH, Na⁺ CP type Iα shRNA (h) Lentiviral Particles: sc-42642-V and Na⁺ CP type Iα shRNA (m) Lentiviral Particles: sc-42643-V.


Molecular Weight of Na⁺ CP type Iα: 260 kDa.

Positive Controls: mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Na⁺ CP type Iα (C-18): sc-16031. Western blot analysis of Na⁺ CP type Iα expression in mouse brain tissue extract.

SELECT PRODUCT CITATIONS

- Basso, K., et al. 2004. Gene expression profiling of hairy cell leukemia reveals a phenotype related to memory B cells with altered expression of chemokine and adhesion receptors. *J. Exp. Med.* 199: 59-68.
- Pinto, F.M., et al. 2009. Molecular and functional characterization of voltage-gated sodium channels in human sperm. *Reprod. Biol. Endocrinol.* 7: 71.