



Jade-1 (E-14): sc-160450

BACKGROUND

Jade-1, also known as PHF17 (PHD finger protein 17), is an 842 amino acid protein that localizes to both the cytoplasm and the nucleus and contains two PHD-type zinc fingers. Expressed at high levels in kidney tissue and also present in heart, pancreas and liver, Jade-1 functions as a transcriptional coactivator that, via its PHD zinc fingers, promotes the TIP30-dependent acetylation of Histone H4. Additionally, Jade-1 is thought to promote apoptosis and is down-regulated in renal carcinoma cells, suggesting a possible role in tumor suppression. Three isoforms of Jade-1, two of which are designated JADE1L and JADE1S, are produced via alternative splicing events. The gene encoding Jade-1 maps to human chromosome 4, which encodes nearly 6% of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all of the human chromosomes.

REFERENCES

- Zhou, M.I., Wang, H., Ross, J.J., Kuzmin, I., Xu, C. and Cohen, H.T. 2002. The von Hippel-Lindau tumor suppressor stabilizes novel plant homeodomain protein Jade-1. *J. Biol. Chem.* 277: 39887-39898.
- Tzouanacou, E., Tweedie, S. and Wilson, V. 2003. Identification of Jade1, a gene encoding a PHD zinc finger protein, in a gene trap mutagenesis screen for genes involved in anteroposterior axis development. *Mol. Cell Biol.* 23: 8553-8552.
- Panchenko, M.V., Zhou, M.I. and Cohen, H.T. 2004. von Hippel-Lindau partner Jade-1 is a transcriptional co-activator associated with histone acetyltransferase activity. *J. Biol. Chem.* 279: 56032-56041.
- Zhou, M.I., Foy, R.L., Chitalia, V.C., Zhao, J., Panchenko, M.V., Wang, H. and Cohen, H.T. 2005. Jade-1, a candidate renal tumor suppressor that promotes apoptosis. *Proc. Natl. Acad. Sci. USA* 102: 11035-11040.
- Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 610514. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Foy, R.L., Song, I.Y., Chitalia, V.C., Cohen, H.T., Saksouk, N., Cayrou, C., Vaziri, C., Côte, J. and Panchenko, M.V. 2008. Role of Jade-1 in the histone acetyltransferase (HAT) HBO1 complex. *J. Biol. Chem.* 283: 28817-28826.
- Chitalia, V.C., Foy, R.L., Bachschmid, M.M., Zeng, L., Panchenko, M.V., Zhou, M.I., Bharti, A., Seldin, D.C., Lecker, S.H., Dominguez, I. and Cohen, H.T. 2008. Jade-1 inhibits Wnt signalling by ubiquitylating β -catenin and mediates Wnt pathway inhibition by pVHL. *Nat. Cell Biol.* 10: 1208-1216.

CHROMOSOMAL LOCATION

Genetic locus: PHF17 (human) mapping to 4q28.2; Phf17 (mouse) mapping to 3 B.

SOURCE

Jade-1 (E-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Jade-1 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-160450 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Jade-1 (E-14) is recommended for detection of Jade-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Jade-1 siRNA (h): sc-88893, Jade-1 siRNA (m): sc-146315, Jade-1 shRNA Plasmid (h): sc-88893-SH, Jade-1 shRNA Plasmid (m): sc-146315-SH, Jade-1 shRNA (h) Lentiviral Particles: sc-88893-V and Jade-1 shRNA (m) Lentiviral Particles: sc-146315-V.

Molecular Weight of Jade-1: 64 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.