

# KIF2B (E-15): sc-160471

## BACKGROUND

The kinesins constitute a large family of microtubule-dependent motor proteins, which are responsible for the distribution of numerous organelles, vesicles and macromolecular complexes throughout the cell. Individual kinesin members play crucial roles in cell division, intracellular transport and membrane trafficking events including endocytosis and transcytosis. KIF2B (kinesin family member 2B) is a 673 amino acid cytoplasmic protein that localizes to the kinetochore. Highly expressed in lung and ovary with moderate expression in heart, kidney, placenta skeletal muscle and spleen, KIF2B has microtubule depolymerization activity and is associated with peripheral translocation of lysosomes. Overexpression of KIF2B in cells result in abnormally large lysosome size and unusual positioning, which is some distance from their usual perinuclear location. KIF2B activity is critical for spindle assembly and chromosome movement.

## REFERENCES

1. Vallee, R.B., Shpetner, H.S. 1990. Motor proteins of cytoplasmic microtubules. *Annu. Rev. Biochem.* 59 : 909-932.
2. Endow, S.A. 1991. The emerging kinesin family of microtubule motor proteins. *Trends Biochem. Sci.* 16: 221-225.
3. Brady, S.T. 1995. A kinesin medley: biochemical and functional heterogeneity. *Trends Cell Biol.* 5: 159-164.
4. Santama, N., et al. 1998. KIF2 $\beta$  a new kinesin superfamily protein in non-neuronal cells, is associated with lysosomes and may be implicated in their centrifugal translocation. *EMBO J.* 17: 5855-5867.
5. Miki, H., et al. 2001. All kinesin superfamily protein, KIF, genes in mouse and human. *Proc. Natl. Acad. Sci. USA* 98: 7004-7011.
6. Homma, N., et al. 2003. Kinesin superfamily protein 2A (KIF2A) functions in suppression of collateral branch extension. *Cell* 114: 229-239.
7. Pfenninger, K.H., et al. 2003. Regulation of membrane expansion at the nerve growth cone. *J. Cell. Sci.* 116: 1209-1217.
8. Manning, A.L., et al. 2007. The kinesin-13 proteins Kif2a, Kif2b, and Kif2c/MCAK have distinct roles during mitosis in human cells. *Mol. Biol. Cell.* 18: 2970-2979.
9. Bakhoum, S.F., et al. 2009. Genome stability is ensured by temporal control of kinetochore-microtubule dynamics. *Nat. Cell Biol.* 11: 27-35.

## CHROMOSOMAL LOCATION

Genetic locus: KIF2B (human) mapping to 17q22.

## SOURCE

KIF2B (E-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of KIF2B of human origin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-160471 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

KIF2B (E-15) is recommended for detection of KIF2B of human and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with KIF2C.

Suitable for use as control antibody for KIF2B siRNA (h): sc-94083, KIF2B shRNA Plasmid (h): sc-94083-SH and KIF2B shRNA (h) Lentiviral Particles: sc-94083-V.

Molecular Weight of KIF2B: 75-80 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.