

Na⁺/K⁺-ATPase α3 (S-19)-R: sc-16051-R

BACKGROUND

The ubiquitously expressed sodium/potassium-ATPase (Na⁺/K⁺-ATPase) exists as a oligomeric plasma membrane complex that couples the hydrolysis of one molecule of ATP to the importation of three Na⁺ ions and two K⁺ ions against their respective electrochemical gradients. As a member of the P-type family of ion motives, Na⁺/K⁺-ATPase plays a critical role in maintaining cellular volume, resting membrane potential and Na⁺-coupled solute transport. Multiple isoforms of three subunits, α, β and γ, comprise the Na⁺/K⁺-ATPase oligomer. The α subunit contains the binding sites for ATP and the cations; the glycosylated β subunit ensures correct folding and membrane insertion of the α subunits. The small γ subunit co-localizes with the α subunit in nephron segments, where it increases the affinity of Na⁺/K⁺-ATPase for ATP. The β subunit, but not the γ subunit, is essential for normal activity of Na⁺/K⁺-ATPase.

REFERENCES

- Hardwicke, P.M., et al. 1981. A proteolipid associated with Na,K-ATPase is not essential for ATPase activity. *Biochem. Biophys. Res. Commun.* 102: 250-257.
- Ackermann, U., et al. 1990. Mutual dependence of Na/K-ATPase α and β subunits for correct posttranslational processing and intracellular transport. *FEBS Lett.* 269: 105-108.
- McDonough, A.A., et al. 1990. The sodium pump needs its β subunit. *FASEB J.* 4: 1598-1605.
- Pedemonte, C.H., et al. 1990. Chemical modification as an approach to elucidation of sodium pump structure-function relations. *Am. J. Physiol.* 258: C1-C23.
- DeTomaso, A.W., et al. 1993. Expression, targeting, and assembly of functional Na/K-ATPase polypeptides in baculovirus-infected insect cells. *J. Biol. Chem.* 268: 1470-1478.
- Mercer, R.W., et al. 1993. Molecular cloning and immunological characterization of the γ polypeptide, a small protein associated with the Na,K-ATPase. *J. Cell Biol.* 121: 579-586.

CHROMOSOMAL LOCATION

Genetic locus: ATP1A3 (human) mapping to 19q13.2; Atp1a3 (mouse) mapping to 7 A3.

SOURCE

Na⁺/K⁺-ATPase α3 (S-19)-R is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of Na⁺/K⁺-ATPase α3 of human origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16051 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Na⁺/K⁺-ATPase α3 (S-19)-R is recommended for detection of Na⁺/K⁺-ATPase α3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Na⁺/K⁺-ATPase α3 (S-19)-R is also recommended for detection of Na⁺/K⁺-ATPase α3 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Na⁺/K⁺-ATPase α3 siRNA (h): sc-36012, Na⁺/K⁺-ATPase α3 siRNA (m): sc-36013, Na⁺/K⁺-ATPase α3 shRNA Plasmid (h): sc-36012-SH, Na⁺/K⁺-ATPase α3 shRNA Plasmid (m): sc-36013-SH, Na⁺/K⁺-ATPase α3 shRNA (h) Lentiviral Particles: sc-36012-V and Na⁺/K⁺-ATPase α3 shRNA (m) Lentiviral Particles: sc-36013-V.


Molecular Weight of Na⁺/K⁺-ATPase α3: 113 kDa.

Positive Controls: Na⁺/K⁺-ATPase α3 (h): 293 Lysate: sc-158752 or rat brain extract: sc-2392.

DATA



Na⁺/K⁺-ATPase α3 (S-19)-R: sc-16051-R. Western blot analysis of Na⁺/K⁺-ATPase α3 expression in non-transfected: sc-110760 (A) and human Na⁺/K⁺-ATPase α3 transfected: sc-158752 (B) 293 whole cell lysates.



Na⁺/K⁺-ATPase α3 (S-19)-R: sc-16051-R. Immunofluorescence staining of formalin-fixed HepG2 cells showing membrane localization.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Na⁺/K⁺-ATPase α3 (H-4): sc-365744** or **Na⁺/K⁺-ATPase α3 (G-6): sc-376967**, our highly recommended monoclonal alternatives to Na⁺/K⁺-ATPase α3 (S-19).