

# Osgep (Y-14): sc-160634

## BACKGROUND

O-sialoglycoprotein endopeptidases cleave the polypeptide backbone of membrane glycoproteins that contain clusters of O-linked sialoglycans. Osgep (O-sialoglycoprotein endopeptidase), also known as GCPL1, is a 335 amino acid protein that is a member of the peptidase M22 family. Osgep specifically cleaves the 31-Arg-I-Asp-32 bond in glycoporphin A, but it does not cleave desialylated glycoproteins, unglycosylated proteins or glycoproteins that are only N-glycosylated. Though ubiquitously expressed at low levels, highest levels of Osgep are found in liver, skeletal muscle and kidney. The gene encoding Osgep maps to human chromosome 14, which houses over 700 genes and comprises nearly 3.5% of the human genome. Chromosome 14 encodes the Presenilin 1 (PSEN1) gene, which is one of the three key genes associated with the development of Alzheimer's disease (AD). The SERPINA1 gene is also located on chromosome 14 and, when defective, leads to the genetic disorder  $\alpha$ 1-antitrypsin deficiency, which is characterized by severe lung complications and liver dysfunction.

## REFERENCES

- Harrison, L., Ascione, A.G., Takiguchi, Y., Wilson, D.M., Chen, D.J. and Demple, B. 1997. Comparison of the promoters of the mouse (APEX) and human (APE) apurinic endonuclease genes. *Mutat. Res.* 385: 159-172.
- Ikeda, S., Ayabe, H., Mori, K., Seki, Y. and Seki, S. 2002. Identification of the functional elements in the bidirectional promoter of the mouse O-sialoglycoprotein endopeptidase and APEX nuclease genes. *Biochem. Biophys. Res. Commun.* 296: 785-791.
- Seki, Y., Ikeda, S., Kiyohara, H., Ayabe, H., Seki, T. and Matsui, H. 2002. Sequencing analysis of a putative human O-sialoglycoprotein endopeptidase gene (OSGEP) and analysis of a bidirectional promoter between the OSGEP and APEX genes. *Gene* 285: 101-108.
- Heilig, R., Eckenberg, R., Petit, J.L., Fonknechten, N., Da Silva, C., Cattolico, L., Levy, M., Barbe, V., de Berardinis, V., Ureta-Vidal, A., Pelletier, E., Vico, V., Anthouard, V., Rowen, L., et al. 2003. The DNA sequence and analysis of human chromosome 14. *Nature* 421: 601-607.
- Godbolt, A.K., Beck, J.A., Collinge, J., Garrard, P., Warren, J.D., Fox, N.C. and Rossor, M.N. 2004. A Presenilin 1 R278I mutation presenting with language impairment. *Neurology* 63: 1702-1704.
- Ng, A.P., Howe Fong, J., Sijin Nin, D., Hirpara, J.L., Asou, N., Chen, C.S., Pervaiz, S. and Khan, M. 2006. Cleavage of misfolded nuclear receptor corepressor confers resistance to unfolded protein response-induced apoptosis. *Cancer Res.* 66: 9903-9912.
- Stolk, J., Seersholm, N. and Kalsheker, N. 2006.  $\alpha$ 1-antitrypsin deficiency: current perspective on research, diagnosis, and management. *Int. J. Chron. Obstruct. Pulmon. Dis.* 1: 151-160.

## CHROMOSOMAL LOCATION

Genetic locus: OSGEP (human) mapping to 14q11.2; Osgep (mouse) mapping to 14 C1.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## SOURCE

Osgep (Y-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Osgep of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-160634 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

Osgep (Y-14) is recommended for detection of Osgep of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with OSGEPL1.


Osgep (Y-14) is also recommended for detection of Osgep in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Osgep siRNA (h): sc-92142, Osgep siRNA (m): sc-151331, Osgep shRNA Plasmid (h): sc-92142-SH, Osgep shRNA Plasmid (m): sc-151331-SH, Osgep shRNA (h) Lentiviral Particles: sc-92142-V and Osgep shRNA (m) Lentiviral Particles: sc-151331-V.


Molecular Weight of Osgep: 36 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Ramos cell lysate: sc-2216 or JAR cell lysate: sc-2276.

## DATA



Osgep (Y-14): sc-160634. Western blot analysis of Osgep expression in Ramos (A), HeLa (B) and JAR (C) whole cell lysates and mouse brain tissue extract (D).



Osgep (Y-14): sc-160634. Western blot analysis of Osgep expression in HeLa (A) and PC-3 (B) whole cell lysates and mouse lymph node tissue extract (C).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

**MONOS**  
Satisfaction  
Guaranteed

Try **Osgep (H-3): sc-393199**, our highly recommended monoclonal alternative to Osgep (Y-14).