

SPACA3 (C-13): sc-160820

BACKGROUND

SPACA3 (sperm acrosome associated 3), also known as sperm lysozyme-like protein 1, lysozyme-like protein 3, lysozyme-like acrosomal sperm-specific secretory protein ALLP-17, cancer/testis antigen 54 (CT54), LYC3, SPRASA or LYZL3, is a 215 amino acid protein that participates in the fusion and adhesion of sperm and egg plasma membrane during fertilization. Identified as a novel cancer/testis antigen in hematologic malignancies, SPACA3 has the ability to elicit B-cell immune responses in patients with cancer and is considered a potential target for immunotherapy. A member of the glycosyl hydrolase 22 family which is expressed in testis, placenta and epididymis, SPACA3 exists as two alternatively spliced isoforms; SPACA3 isoform 1 is a single-pass type II membrane protein of the sperm acrosome whereas SPACA3 isoform 2 is a secreted protein.

REFERENCES

1. Welsh, I.R. and Spitznagel, J.K. 1971. Distribution of lysosomal enzymes, cationic proteins, and bactericidal substances in subcellular fractions of human polymorphonuclear leukocytes. *Infect. Immun.* 4: 97-102.
2. Cramer, E.M. and Breton-Gorius, J. 1987. Ultrastructural localization of lysozyme in human neutrophils by immunogold. *J. Leukoc. Biol.* 41: 242-247.
3. Ibrahim, H.R., Matsuzaki, T. and Aoki, T. 2001. Genetic evidence that antibacterial activity of lysozyme is independent of its catalytic function. *FEBS Lett.* 506: 27-32.
4. Mandal, A., Klotz, K.L., Shetty, J., Jayes, F.L., Wolkowicz, M.J., Bolling, L.C., Coonrod, S.A., Black, M.B., Diekman, A.B., Haystead, T.A., Flickinger, C.J. and Herr, J.C. 2003. SLLP1, a unique, intra-acrosomal, non-bacteriolytic, c lysozyme-like protein of human spermatozoa. *Biol. Reprod.* 68: 1525-1537.
5. Zhang, K., Gao, R., Zhang, H., Cai, X., Shen, C., Wu, C., Zhao, S. and Yu, L. 2005. Molecular cloning and characterization of three novel lysozyme-like genes, predominantly expressed in the male reproductive system of humans, belonging to the c-type lysozyme/ α -lactalbumin family. *Biol. Reprod.* 73: 1064-1071.
6. Niyonsaba, F. and Ogawa, H. 2005. Protective roles of the skin against infection: implication of naturally occurring human antimicrobial agents β -defensins, cathelicidin LL-37 and lysozyme. *J. Dermatol. Sci.* 40: 157-168.

CHROMOSOMAL LOCATION

Genetic locus: SPACA3 (human) mapping to 17q11.2.

SOURCE

SPACA3 (C-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal extracellular domain of SPACA3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-160820 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SPACA3 (C-13) is recommended for detection of SPACA3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other SPACA family members.

SPACA3 (C-13) is also recommended for detection of SPACA3 in additional species, including bovine.

Suitable for use as control antibody for SPACA3 siRNA (h): sc-93867, SPACA3 shRNA Plasmid (h): sc-93867-SH and SPACA3 shRNA (h) Lentiviral Particles: sc-93867-V.

Molecular Weight of SPACA3: 23 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.